

# Age-Related Changes In Female and Male Cerebral Volumes

## A Study of Whole-Brain Tissue Using the Stereological Method

Remzi Yigiter\*, Erol Avşın<sup>†</sup>, Atakan Yücel<sup>‡</sup>, Adem Kara<sup>§</sup>, Nermin Yücel<sup>||</sup>,  
Ünsal Aydınöğlü<sup>‡</sup>, İsmail Can<sup>§</sup>, Mustafa Güleç<sup>‡</sup>, Samih Diyarbakır<sup>†</sup>

### ABSTRACT

The stereological techniques based on the Cavalieri methods are used in many studies to estimate volumes of anatomical structures in an unbiased fashion. This is a rapid, inexpensive approach that provides a correct volumetric calculation using magnetic resonance (MR) brain images. We investigated and age-sex-related volumetric changes in human brain using Cavalieri method. Three-dimensional MR imaging was obtained 120 healthy volunteers to 10 people from each age and gender-decade group. The Cavalieri method of modern design stereology was used to measure the cerebral volumes. To estimate this volume, the point counting approach was preferred. In our study, it was found that significant decreases in whole-brain volume ( $p < 0.05$ ) emerge progressively with increasing age. The most remarkable alteration was observed in older people's brain volume. There was significant volumetric effect of gender on normalized brain volumes. Furthermore, both women and men showed statistically significant asymmetries in hemisphere volume. This quantitative volumetric study demonstrated a possible normal increase and then decrease in brain volume from age 0 to 60 years in healthy subjects. The differences in the brain sizes between the male and female subjects may be mainly attributable to the differences in total brain volume. A better understanding of this process may help to distinguish normal age-related alterations from neurodegenerative diseases.

**Key Words:** brain volume, stereology, MR images, age-sex-related volumetric changes, developmental change

NeuroQuantology 2012; 1: 132-139

### 1. Introduction

The quantitative assessment of brain volume is an increasingly important consideration in clinical studies and the treatment of diseases such as Alzheimer's, multiple sclerosis, and schizophrenia. For this reason, magnetic resonance imaging (MRI) technology has been used to perform morphometric studies on the brain (Odaci *et al.*, 2003; Elfaki *et al.*, 2011). Age-related changes in brain volume in healthy adult brains have also been of great

interest in recent years, because the determination of normal age-specific brain volume values is very important when it comes to evaluating both pathological conditions and the normal aging process (Biegonet *et al.*, 1994). The estimated results from such analyses have shown that age-related brain loss may vary in different brain regions, as well as according to hemisphere (Miller *et al.*, 1980; Coffey *et al.*, 1992) and sex (Xu *et al.*, 2000). Human cerebral volume estimation via stereological methods is based on MRI. One of these methods is Cavalieri method, which is a sampling technique used to generate mathematically unbiased estimates of the geometric objects of three-dimensional structures based on two-dimensional slices of the object. One of the most important quantitative parameters is volume, which represents important data in many research areas (Unal *et al.*, 2010a). The stereological method has been mathematically validated, and this method should be used to determine

**Corresponding author:** Atakan Yücel, MD., Office address: Erzurum Ataturk University, Faculty of Medicine, Departments of Psychiatry, Erzurum, Turkey.

**Address:** \*Departments of Neurology, Medical Faculty of Gaziantep University, Gaziantep; Turkey. <sup>†</sup>Departments of Anatomy, Medical Faculty of Ataturk University, Erzurum; Turkey. <sup>‡</sup>Departments of Psychiatry, Medical Faculty of Ataturk University, Erzurum; Turkey. <sup>§</sup>Departments of Histology, Medical Faculty of Ataturk University, Erzurum; Turkey. <sup>||</sup>Departments of Child Psychiatry, Medical Faculty of Ataturk University, Erzurum; Turkey.

**Phone:** +904422317410

✉ dr\_atakanyucel@hotmail.com

Received Sept 29, 2011. Revised Dec 1, 2011. Accepted Jan 6, 2012.

eISSN 1303-5150



the volumetric effects of histopathologic materials.

In humans, normal aging can affect brain volumes in as life progresses (Drayer, 1988). T2-weighted MRI measured brain volume decreases with increasing age, while sex differences affect human brain volume during progressive life; a decrease in brain volume correlates well with impaired cognitive function (Coffey *et al.*, 1998). However, the structural alterations that cause the development and progression of brain atrophy are not yet fully understood (Good *et al.*, 2001). Many studies have suggested that cerebral hemispheric symmetry or asymmetry of brain volume is related with an interaction between brain function and morphology (Watkins *et al.*, 2001). The aim of the present study was to investigate possible brain volume changes via stereological methods in normal adults and adolescents divided into 23 groups according to age and gender using MRI. In addition, we also compared age-dependent brain volumes in both hemispheres.

## 2. Materials and Methods

### MRI

A total of 120 subjects who visited the hospital for a personal health-screening program participated in this study. MRI (1.5-T MRI unit, Magnetom Impact, Siemens Medical Systems) was obtained for 10 healthy volunteer visitors in each group (decade 1: age 0–10, decade 2: age 10–20; decade 3: age 20–30; decade 4: 30–40 age, decade 5: 50–50; and decade 6: 50–60). We used the turbo spin echo technique (TR 4,600 ms, TE 90 ms, FOV 201 x 230 mm, matrix 132 x 256.3 mm thick slices with a 1-2 mm gap, and three acquisitions) to obtain MRI whole-brain images, which thickness was approximately 5-6 mm. In this way, we obtained 15–16 images for each patient's whole brain. All measurements and estimations were performed on these images (Fig. 1).

### MRI Stereological Estimations

We used the Cavalieri method in modern design-based stereology to measure cerebral volumes. To perform the estimation, the point counting approach was preferred. Firstly, detailed systematic series of axial MRI (T1-weighted, TR/TE 400/10 msec) of 5 mm thickness were obtained throughout the whole brain of subjects. Secondly, to magnify the

images, all MRI were projected on a screen using a projection machine, and then the images were evaluated with the point counting method (Gundersen, 1986; Gundersen *et al.*, 1988; Cruz-Orive and Weibel, 1990). We used the serial and parallel section techniques with a known distance of these images to estimate the considered objects' volumes correctly; this method is first important rule for stereology (Gundersen, 1986; Gundersen *et al.*, 1988; Cruz-Orive and Weibel, 1990). We performed a pilot study before beginning the study to set both the coefficient of error (CE) and variation, as well as to modify our approach using the imaging technique according to stereological principles and determine the slice thickness from considered object. We applied the systematic Random Sampling, while the samples were chosen and all interest object images were evaluated using stereological methods (Mackay *et al.*, 1999).

Images obtained from the objects considered were projected onto a screen using a projector (Braun Paxiscope 650) (Figure 1). A point grid was employed for cerebral volume estimations, and the grid points were equally spaced (0.5 cm) both across and down. The grid point density was determined as a serial MRI number, which was taken as the CE. The CE was calculated according to formula of Gundersen and Jensen, and 13–15 cerebral images were used to measure cerebral volumes.

The point grid was randomly placed on the MRI to measure the area of interest and count how many points hit the region of interest according to the constant rule. The results were then used to compute the volume (Fig. 2).





Figure 1. Images of interest were projected onto a screen using a projector





**Figure 2.** Estimation of the area of the slice using a transparent point grid

The test point of grid area was  $a \text{ cm}^2$  and the size was  $k \text{ cm}^2$  ( $a \text{ cm}^2 = k \times k \text{ cm}^2$ ). We calculated the MRI and overhead projector magnifications, and then multiplied by the magnification of the calculated values ( $\text{cm}^2$ ):

$$\text{Volume} = t \text{ (cm)} \times a/p \text{ (cm} \times \text{cm)} \times ((\sum P) \text{ cm}^3)$$

Here,  $V$  is the volume of the object of interest object in one section plane,  $t$  is the section thickness,  $a/p$  is the inter point area, and  $\sum P$  is the number of points touching the area of interest in that section (Gundersen *et al.*, 1988). The volumes for each section were estimated by the above formula and the sum of total brain volumes was estimated by the formula given below:

$$\text{Total Volume} = V_1 + V_2 + \dots + V_n$$

The CE of the volume was estimated due to the combined effects of sectioning and point counting (Gundersen and Jensen, 1987). The formula for CE estimation is as follows:

$$\text{Noise} = 0.0724 \times (b / \sqrt{a}) \times \sqrt{n \times \sum P}$$

Here, Noise represents the sample section near-surface reflection data,  $b / \sqrt{a}$  denotes the average length of the boundary divided by the square root of the average area of the boundary,  $n$  is the number of slices of interest, and  $\sum P$  is the total number of points hitting the object of interest.

$$\text{Var}_{\text{SRS}} \left( \sum_{i=1}^n a \right) = (3 \cdot (A - \text{Noise}) - 4 \cdot B + C) / 12$$

SRS in this equation stands for systematic random sampling. Here,

$$\text{Var}_{\text{SRS}} \left( \sum_{i=1}^n a \right)$$

is the variation in total area. These data gave us information about what number of slices would be sufficient. A, B, and C are shown in the columns in Table 1.

$$\text{TotalVar} = \text{Noise} + \text{Var}_{\text{SRS}}$$

$$\text{CE}(\sum P) = \frac{\sqrt{\text{Total Var}}}{\sum P}$$

$$\begin{aligned} \text{Noise} &= 0.0724 \times (b / \sqrt{a}) \times \sqrt{n \times \sum P} \\ &= 0.0724 \times 5 \times \sqrt{227 \times 10} = 17.25 \end{aligned}$$

$$\begin{aligned} \text{Var}_{\text{SRS}} \left( \sum_{i=1}^n a \right) &= (3 \times (A - \text{Noise}) - 4 \times B + C) / 12 \\ &= (3 \times (5251 - 17.25) - 4 \times 4783 + 4371) / 12 = 78.35 \end{aligned}$$

$$\text{Total Var} = \text{Noise} + \text{Var}_{\text{SRS}} = 17.25 + 78.35 = 95.6$$

$$\text{CE}(\sum P) = \frac{\sqrt{\text{Total Var}}}{\sum P} = \frac{\sqrt{95.6}}{227} = 0.043$$

$$\begin{aligned} \text{Volume} &= t \times a/p \times ((\sum P) - (1/2)) \\ &= 1 \times 3.642 \times (289.3 - 16) = 995.48 \text{ cm}^3 \end{aligned}$$

### Statistical Analyses

For statistical analysis, differences between the decades were tested by the analysis of variance (ANOVA) and paired samples test using SPSS 17.0 for Windows XP (SPSS Inc., Chicago, IL). A value of  $p < 0.05$  was considered significant.

### 3. Results

In this study, the brain volumes of healthy men and women in all age-decade groups were shown in Table 2.

Firstly, the average cerebral volumes of subjects' of the same age and gender were compared with those of older groups of the same gender for all age-decade groups. The same comparison was applied to both sexes (Table 3). With the exception of the last comparison for males, the pairs of comparisons showed a statistically significant difference ( $p < 0.05$ ).

Secondly, age-dependent average cerebral volume in the same group was compared by gender. The average brain volume difference was 6.36% for males; all pairs of comparisons were shown in Table 4.



When the right and left hemisphere volumes were compared in groups from the same decade, the pair of group comparisons and the results were seen in Table 5. When the male and female right and left hemispheres volumes were compared according to gender, all comparisons except males in decade 6 and females in decade 1 showed a statistically significant difference; the comparisons were seen in Table 5.

**Table 1.** In this study, estimates of brain volumes were generated according to the sampling strategy procedure.

Section No	P <sub>i</sub>	P <sub>i</sub> x P <sub>i</sub>	P <sub>i</sub> x P <sub>i+1</sub>	P <sub>i</sub> x P <sub>i+2</sub>
1	20	400	440	460
2	22	484	506	528
3	23	529	552	667
4	24	576	696	600
5	29	841	725	725
6	25	625	625	525
7	25	625	525	425
8	21	441	357	441
9	17	289	357	0
10	21	441	0	0
	<b>227</b>	<b>5251</b>	<b>4783</b>	<b>4371</b>

**Table 2.** Male and female average brain volumes.

	Female (N:10)	Male (N: 10)
1 <sup>st</sup> decade	1404.358 cm <sup>3</sup>	1415.296 cm <sup>3</sup>
2 <sup>nd</sup> decade	1592.826 cm <sup>3</sup>	1613.766 cm <sup>3</sup>
3 <sup>th</sup> decade	1650.087 cm <sup>3</sup>	1769.144 cm <sup>3</sup>
4 <sup>th</sup> decade	1423.788 cm <sup>3</sup>	1505.39 cm <sup>3</sup>
5 <sup>th</sup> decade	1282.099 cm <sup>3</sup>	1452.987 cm <sup>3</sup>
6 <sup>th</sup> decade	1267.135 cm <sup>3</sup>	1409.906 cm <sup>3</sup>

**Table 3.** Comparison of female and male cerebral volume in older groups.

Group	Difference One Way ANOVA
1 <sup>st</sup> decade / 2 <sup>nd</sup> decade female	(P<0.05)
2 <sup>nd</sup> decade / 3 <sup>th</sup> decade female	(P<0.05)
3 <sup>th</sup> decade / 4 <sup>th</sup> decade female	(P<0.05)
4 <sup>th</sup> decade / 5 <sup>th</sup> decade female	(P<0.05)
5 <sup>th</sup> decade / 6 <sup>th</sup> decade female	(P>0.05)
1 <sup>st</sup> decade / 2 <sup>nd</sup> decade male	(P<0.05)
2 <sup>nd</sup> decade / 3 <sup>th</sup> decade male	(P<0.05)
3 <sup>th</sup> decade / 4 <sup>th</sup> decade male	(P<0.05)
4 <sup>th</sup> decade / 5 <sup>th</sup> decade male	(P<0.05)
5 <sup>th</sup> decade / 6 <sup>th</sup> decade male	(P>0.05)

In this study, we also investigated the difference between the total cerebral right hemisphere volumes by gender; the difference was 8.20% in favor of the male right hemisphere. The groups comparisons are

shown in Table 6. Similar results were obtained from comparisons of the total cerebral left hemisphere volumes by gender; the difference was 3.68% in favor of the male left hemisphere. The group comparisons are shown in Table 7.

**Table 4.** Comparison of the same age groups' total brain volumes by sex.

Group	Difference – One way ANOVA, all p value
1 <sup>st</sup> decade male/ 1 <sup>st</sup> decade female	0.06% (on behalf of male. P>0.05)
2 <sup>nd</sup> decade male/ 2 <sup>nd</sup> decade female	1.30% (on behalf of male. P<0.05)
3 <sup>th</sup> decade male/ 3 <sup>th</sup> decade female	7.23% (on behalf of male. P<0.05)
4 <sup>th</sup> decade male/ 4 <sup>th</sup> decade female	5.74% (on behalf of male. P<0.05)
5 <sup>th</sup> decade male/ 5 <sup>th</sup> decade female	13.35% (on behalf of male. P<0.05)
6 <sup>th</sup> decade male / 6 <sup>th</sup> decade female	11.44% (on behalf of male. P<0.05)

When female subjects' age-dependent cerebral left hemisphere volumes were compared to those of the older female groups, all groups were statistically significantly different from one another (one-way ANOVA, p<0.05). The group comparisons are shown in Table 8. When female age-dependent cerebral right hemisphere volumes were compared to the older female groups, all groups comparisons except decade 5/decade 6 comparisons were statistically significantly different from each other (one-way ANOVA, p<0.05). The group comparisons are shown in Table 9.

In the same way, male cerebral left hemispheres were compared to those of the older male groups. All group comparisons except decade 5/decade 6 were statistically significantly different from the other group comparisons (one-way ANOVA, p<0.05). The group comparisons are shown in Table 10. When male subjects' right hemispheres were compared to those of males in the older groups, all comparisons except decade 5/decade 6 were statistically significantly different (one-way ANOVA, p<0.05). The group comparisons are shown in Table 11.





**Table 5.** Comparison of male and female cerebral volumes by hemisphere volumes.

Group	Left	Right	Difference
1 <sup>st</sup> decade female left/right	701.199 cm <sup>3</sup>	703.159 cm <sup>3</sup>	0.28% (on behalf of right hemisphere. P>0.05)
2 <sup>nd</sup> decade female left/right	817.745 cm <sup>3</sup>	775.081 cm <sup>3</sup>	5.28% (on behalf of left hemisphere. P<0.05)
3 <sup>th</sup> decade female left/right	833.285 cm <sup>3</sup>	816.802 cm <sup>3</sup>	1.98% (on behalf of left hemisphere. P<0.05)
4 <sup>th</sup> decade female left/right	721.605 cm <sup>3</sup>	702.183 cm <sup>3</sup>	2.7% (on behalf of left hemisphere. Test. P<0.05)
5 <sup>th</sup> decade female left/right	653.991 cm <sup>3</sup>	628.108 cm <sup>3</sup>	3.96% (on behalf of left hemisphere. P<0.05)
6 <sup>th</sup> decade female left/right	641.456 cm <sup>3</sup>	625.679 cm <sup>3</sup>	2.46% (on behalf of left hemisphere. P<0.05)
1 <sup>st</sup> decade male left/right	699.651cm <sup>3</sup>	715.645 cm <sup>3</sup>	2.28% (on behalf of right hemisphere. P<0.05)
2 <sup>nd</sup> decade male left/right	797.374cm <sup>3</sup>	816.392 cm <sup>3</sup>	2.38% (on behalf of right hemisphere. P<0.05)
3 <sup>th</sup> decade male left/right	876.575cm <sup>3</sup>	892.569 cm <sup>3</sup>	1.82% (on behalf of right hemisphere. P<0.05)
4 <sup>th</sup> decade male left/right	743.757 cm <sup>3</sup>	761.633 cm <sup>3</sup>	2.4% (on behalf of right hemisphere. P<0.05)
5 <sup>th</sup> decade male left/right	718.944 cm <sup>3</sup>	734.043 cm <sup>3</sup>	2.1% (on behalf of right hemisphere. P<0.05)
6 <sup>th</sup> decade male left/right	699.767 cm <sup>3</sup>	710.139 cm <sup>3</sup>	1.48% (on behalf of right hemisphere. P>0.05)

**Table 6.** Comparison of the same age groups' right hemisphere by sex.

Group	Difference
Paired Samples Test. One way ANOVA	
1 <sup>st</sup> decade male right/female right brain volume	1.75% (on behalf of male right hemisphere. P<0.05)
2 <sup>nd</sup> decade male right/female right brain volume	5.07% (on behalf of male right hemisphere. P<0.05)
3 <sup>th</sup> decade male right/female right brain volume	8.49% (on behalf of male right hemisphere. P<0.05)
4 <sup>th</sup> decade male right/female right brain volume	7.81% (on behalf of male right hemisphere. P<0.05)
5 <sup>th</sup> decade male right/female right brain volume	14.44% (on behalf of male right hemisphere. P<0.05)
6 <sup>th</sup> decade male right/female right brain volume	11.9% (on behalf of male right hemisphere. P<0.05)

**Table 7.** Comparison of the same age groups' left hemisphere by sex.

Group	Difference
Paired Samples Test. One way ANOVA	
1 <sup>st</sup> decade female left /1st decade male left brain volume	0.22% (decreased left hemisphere volume on behalf of female. P>0.05)
2 <sup>nd</sup> decade female left / 2 <sup>nd</sup> decade male left brain volume	2.55% (decreased left hemisphere volume on behalf of female. P<0.05)
3 <sup>th</sup> decade female left /3 <sup>th</sup> decade male left brain volume	4.94% (increased left hemisphere volume on behalf of male. P<0.05)
4 <sup>th</sup> decade female left /4 <sup>th</sup> decade male left brain volume	2.98% (increased left hemisphere volume on behalf of male. P<0.05)
5 <sup>th</sup> decade female left /5 <sup>th</sup> decade male left brain volume	9.94% (increased left hemisphere volume on behalf of male. P<0.05)
6 <sup>th</sup> decade female left /6 <sup>th</sup> decade male left brain volume	8.34% (on behalf of male left hemisphere. P<0.05)

**Table 8.** Comparison of female left cerebral hemispheres in older groups.

Group	Difference
1 <sup>st</sup> decade female left/2 <sup>nd</sup> decade female left brain volume	16.62% (increased volume on behalf of older age group. P<0.05)
2 <sup>nd</sup> decade female left/3 <sup>th</sup> decade age female left brain volume	1.9% (increased volume on behalf of older age group P<0.05)
3 <sup>th</sup> decade female left/4 <sup>th</sup> decade female left brain volume	13.41% (decreased volume on behalf of older age group. P<0.05).
4 <sup>th</sup> decade female left/5 <sup>th</sup> decade female left brain volume	9.37%(decreased volume on behalf of older age group P<0.05).
5 <sup>th</sup> decade female left/6 <sup>th</sup> decade female left brain volume	1.92% (decreased volume on behalf of older age group. P<0.05)

**Table 9.** Comparison of female right cerebral hemispheres in older groups.

Group	Difference
1 <sup>st</sup> decade female right/2 <sup>nd</sup> decade female right brain volume	10.23% (increased volume on behalf of older age group. P<0.05)
2 <sup>nd</sup> decade female right/3 <sup>th</sup> decade female right brain volume	5.38% (increased volume on behalf of older age group. P<0.05)
3 <sup>th</sup> decade female right/4 <sup>th</sup> decade female right brain volume	16.32% (decreased volume on behalf of older age group. P<0.05).
4 <sup>th</sup> decade female right/5 <sup>th</sup> decade female right brain volume	11.20% (decreased volume on behalf of older age group. P<0.05).
5 <sup>th</sup> decade female right/6 <sup>th</sup> decade female right brain volume	0.49% (decreased volume on behalf of older age group. P>0.05)



**Table 10.** Comparison of male left cerebral hemispheres in older groups.

Group	Difference
1 <sup>st</sup> decade male left/2 <sup>nd</sup> decade male left brain volume	12.26% (increased volume on behalf of older age group. P<0.05)
2 <sup>nd</sup> decade left/3 <sup>th</sup> decade male left brain volume	9.04%(increased volume on behalf of older age group. P<0.05)
3 <sup>th</sup> decade male left/4 <sup>th</sup> decade left brain volume	17.85% (decreased volume on behalf of older age group. P<0.05).
4 <sup>th</sup> decade male left/5 <sup>th</sup> decade male left brain volume	3.45% (decreased volume on behalf of older age group. P<0.05).
5 <sup>th</sup> decade male left/6 <sup>th</sup> decade male left brain volume	2.75% (decreased volume on behalf of older age group. P>0.05)

**Table 11.** Comparison of male right cerebral hemispheres in older groups.

Group	Difference, One way ANOVA
1 <sup>st</sup> decade male/2 <sup>nd</sup> decade male right brain volume	14.07% (increased volume on behalf of older group. P<0.05)
2 <sup>nd</sup> decade male/3 <sup>th</sup> decade male right brain volume	9.33% (increased volume on behalf of older group. P<0.05)
3 <sup>th</sup> decade male /4 <sup>th</sup> decade male right brain volume	14.66% (decreased volume on behalf of older group. P<0.05)
4 <sup>th</sup> decade male /5 <sup>th</sup> decade male right brain volume	3.62% (decreased volume on behalf of older group. P<0.05)
5 <sup>th</sup> decade male /6 <sup>th</sup> decade male right brain volume	3.25% (decreased volume on behalf of older group. P>0.05)

#### 4. Discussion

In this study, we measured both male and female cerebral volumes in healthy subjects aged 0–60 years and evaluated the data by age, sex, and hemisphere. The results showed patterns of age-related changes in male and female brain volumes via the Cavalieri method (Table 2). The female brain volume results can be summarized as follows: A regular and constant increase was observed in cerebral volumes from decade 1 to decade 4, but cerebral volume started to decrease from decade 4 to decade 6. All group differences were statistically significant ( $p < 0.05$ ). In male subjects, brain volumes constantly increased in decade 1 to decade 3, but decreased from decade 4 to decade 6. Except for decade 5/decade 6, all group comparisons exhibited statistical significance ( $p < 0.05$ ) (Table 3). In some research, while brain volume and weight have been found to change according to age and gender, brain ventricles are expanded (Anderton, 1997; Anderton, 2002). We found similar results in our study, but the decade 5 and decade 6 groups exhibited decreased brain volumes. In addition, the brain volumes

increased to decade 3, then decreased by small amounts ( $p > 0.05$ ).

Some researchers have tried to clarify whether neuronal pigmentation and granulo-vascular degeneration are related to brain volume changes (Ball, 1977). One stereological study found that total brain volume was increased by 15% and myelination nervous fibrils were increased by 17% in adolescents when compared to adult subjects (Tang *et al.*, 1997). The findings from the study showed a linear decrease of 0.31% of total brain volume per year in subjects at an average age of 73 years (Ikram *et al.*, 2008). A parallel decrease of 0.3% per year was reported in the population-based WHICAP cohort of people aged an average of 80 years (Brickman *et al.*, 2008). Some related studies have shown that the myelination of nervous fibrils decreased at 30% with increasing age, but this alteration showed differences between sexes (Marnier *et al.*, 2003). Sex-dependent degenerative changes affected female subjects less than male subjects; the reason for this is that female gonadal hormones have a neuroprotective effect on the nervous system (Unal *et al.*, 2010b). Our study results are in accordance with these findings.

One stereological study reported that investigated the cerebrum (cortex, basal ganglions, thalamus, and white matter), cerebellum, and lateral ventricle volumes of 32 female and 32 male subjects aged 74–87 over four years. It was shown that cerebral volumes decreased 2.1% per year, lateral ventricles volumes expanded 5.6% per year, and the progression of cerebral atrophy was statistically significant by sex (Tang *et al.*, 2001). In our study, brain volumes showed a linear increase with age in adolescents, and then a decrease that was stronger in men than in women. The linear decrease of brain volume has also reported in other studies (DeCarli *et al.*, 2005).

Researchers have been trying to clarify the differences between the male and female cortical functions for decades. Therefore, the present study investigated the relationship between hemispheric asymmetries and morphological changes between sexes (Harshman and Remington, 1976; Shaywitz *et al.*, 1995; Hirnstein *et al.*, 2010). The relationship between hemispheres linearly is positive, because hemispheric asymmetries originate in better cognitive processing.



However, the experimental data supporting this concept are mixed, and the statistical methods to analyze this relationship have been assessed. When the brain's morphological structure was evaluated in relation to sexual dimorphism, sexual dimorphism was found in the corpus callosum, or splenium, and this asymmetry appeared in the decade 4 and older age groups (Dubb *et al.*, 2003); the male brain volumes decreased to a lesser extent than those of females in adult subjects (Gur *et al.*, 2002). These findings showed that degenerative changes varied by sex (Kertesz *et al.*, 1990). Sex-dependent degenerative changes did not appear in early adolescence, however, and the study indicated that male and female brain volumes begin to change at the age of 10 years (Giedd *et al.*, 1997). The brain volume changes between the sexes observed in the present study might be the result of hormonal activity in the body. In our study, we found similar results, where in the male and female subjects' brain volumes started to change at 10 years old. Compared to decade 1 total brain volumes by sex, no statistically significant difference was determined in groups between the sexes. However, the male and female brain volume difference increased in relation to increasing age.

Age-dependent male and female brain volumes have been reported to be different from each other, and male brain volumes have been found to be higher than female brain volumes in many studies (Giedd *et al.*, 1997). This finding is in accordance with our results, where in male and female brain volumes

exhibited a statistically significant difference in favors of male.

As far as hemispheric asymmetries are concerned, according to our method, the brain showed volumetric differences by hemisphere as well as sex. While male right hemisphere volumes were higher than left hemisphere volumes, female left hemisphere volumes were higher than right hemisphere volumes. Although some studies support this notion (Gur *et al.*, 1991), others have reported results that are contrary to our findings (Sullivan *et al.*, 2001; Gur *et al.*, 2002).

## 5. Conclusion and Outlook

This quantitative volumetric study demonstrated a possible normal increase followed by a decrease of brain volume across adult life, from 0–60 years, in healthy subjects. The volume changes in different decades indicate that brain volume increase occurs in a linear pattern starting from early adolescence (0–10 years), whereas brain volume decrease occurs in early adulthood (30–40 years). These alterations illustrate processes of both brain maturation and aging during adult and adolescent life, and they may be taken as targets in evaluating pathological conditions such as atrophy with increasing age. Differences in brain sizes between male and female subjects may be mainly attributable to differences in total brain volume. A better understanding of this process may help to distinguish normal age-related alterations from neurodegenerative diseases.



## References

- Anderton BH. Changes in the ageing brain in health and disease. *Philos Trans R Soc Lond B Biol Sci* 1997; 352: 1781-1792.
- Anderton BH. Ageing of the brain. *Mech Ageing Dev* 2002; 123: 811-817.
- Ball MJ. Neuronal loss, neurofibrillary tangles and granulovacuolar degeneration in the hippocampus with ageing and dementia. A quantitative study. *Acta Neuropathol* 1977; 37: 111-118.
- Biegona A, Eberling JL, Richardson BC, Roos MS, Wong ST, Reed BR, Jagust WJ. Human corpus callosum in aging and Alzheimer's disease: a magnetic resonance imaging study. *Neurobiol Aging* 1994; 15: 393-397.
- Brickman AM, Schupf N, Manly JJ, Luchsinger JA, Andrews H, Tang MX, Reitz C, Small SA, Mayeux R, DeCarli C, Brown TR. Brain morphology in older African Americans, Caribbean Hispanics, and whites from northern Manhattan. *Arch Neurol* 2008; 65: 1053-1061.
- Coffey CE, Wilkinson WE, Parashos IA, Soady SA, Sullivan RJ, Patterson LJ, Figiel GS, Webb MC, Spritzer CE, Djang WT. Quantitative cerebral anatomy of the aging human brain: a cross-sectional study using magnetic resonance imaging. *Neurology* 1992; 42: 527-536.
- Coffey CE, Lucke JF, Saxton JA, Ratcliff G, Unitas LJ, Billig B, Bryan RN. Sex differences in brain aging: a quantitative magnetic resonance imaging study. *Arch Neurol* 1998; 55: 169-179.
- Cruz-Orive LM, Weibel ER. Recent stereological methods for cell biology: a brief survey. *Am J Physiol* 1990; 258: 148-156.
- DeCarli C, Massaro J, Harvey D, Hald J, Tullberg M, Au R, Beiser A, D'Agostino R, Wolf PA. Measures of brain morphology and infarction in the Framingham heart study: establishing what is normal. *Neurobiol Aging* 2005; 26: 491-510.
- Drayer BP. Imaging of the aging brain. Part I. Normal findings. *Radiology* 1988; 166: 785-796.
- Dubb A, Gur R, Avants B, Gee J. Characterization of sexual dimorphism in the human corpus callosum. *Neuroimage* 2003; 20: 512-519.
- Elfaki A, Osman T, Elsheikh A, Hamdoun A, Sahin B. Evaluation of the intra-rater variation for the estimation of volume of cerebral structures using the cavalieri principle on magnetic resonance images. *J Exp Clin Med* 2011; 28:22-25.
- Giedd JN, Castellanos FX, Rajapakse JC, Vaituzis AC, Rapoport JL. Sexual dimorphism of the developing human brain. *Prog Neuropsychopharmacol Biol Psychiatry* 1997; 21: 1185-1201.
- Good CD, Johnsrude IS, Ashburner J, Henson RN, Friston KJ, Frackowiak RS. A voxel-based morphometric study of ageing in 465 normal adult human brains. *Neuroimage* 2001; 14: 21-36.
- Gur RC, Mozley PD, Resnick SM, Gottlieb GL, Kohni M, Zimmerman R, Herman G, Atlas S, Grossman R, Berretta D. Gender differences in age effect on brain atrophy measured by magnetic resonance imaging. *Proc Natl Acad Sci USA* 1991; 88: 2845-2849.
- Gur RC, Gunning-Dixon FM, Turetsky BI, Bilker WB, Gur RE. Brain region and sex differences in age association with brain volume: a quantitative MRI study of healthy young adults. *Am J Geriatr Psychiatry* 2002; 10: 72-80.
- Gundersen HJ, Jensen EB. The efficiency of systematic sampling in stereology and its prediction. *J Microsc* 1987; 147: 229-263.
- Gundersen HJ. Stereology of arbitrary particles. A review of unbiased number and size estimators and the presentation of some new ones, in memory of William R. Thompson. *J Microsc* 1986; 143: 3-45.
- Gundersen HJG, Bendtsen TF, Korbo L, Marcussen N, Møller A, Nielsen K, Nyengaard JR, Pakkenberg B, Sørensen FB, Vesterby Aand West MJ. Some new, simple and efficient stereological methods and their use in pathological research and diagnosis. *APMIS* 1988; 96: 379-394.
- Hirnstain M, Leask S, Rose J, Hausmann M. Disentangling the relationship between hemispheric asymmetry and cognitive performance. *Brain Cogn* 2010; 73: 119-127.
- Harshman R.A. and Remington R. Sex, language and the brain: a review of the literature on adult sex differences in lateralization. *UCLA Working Papers Phonet* 1976; 31: 86-103.
- Ikram MA, Vrooman HA, Vernooij MW, van der Lijn F, Hofman A, van der Lugt A, Niessen WJ, Breteler MM. Brain tissue volumes in the general elderly population. The Rotterdam Scan Study. *Neurobiol Aging* 2008; 29: 882-890.
- Kertesz A, Polk M, Black SE, Howell J. Sex, handedness and the morphometry of cerebral asymmetries on magnetic resonance imaging. *Brain Res* 1990; 530: 40-48.
- Mackay CE, Pakkenberg B, Roberts N. Comparison of compartment volumes estimated from MR images and physical sections of formalin fixed cerebral hemispheres. *Acta Stereol* 1999; 18: 149-159.
- Marnier L, Nyengaard JR, Tang Y, Pakkenberg B. Marked loss of myelinated nerve fibers in the human brain with age. *J Comp Neurol* 2003; 462: 144-152.
- Miller AK, Alston RL, Corsellis JA. Variation with age in the volumes of grey and white matter in the cerebral hemispheres of man: measurements with an image analyser. *Neuropath Appl Neuro* 1980; 6: 119-132.
- Odaci E, Sahin B, Sonmez OF, Kaplan S, Bas O, Bilgic S, Bek Y, Ergür H. Rapid estimation of the vertebral body volume, a combination of the Cavalieri principle and computed tomography images. *Eur J Radiol* 2003; 48: 316-326.
- Shaywitz BA, Shaywitz SE, Pugh KR, Constable RT, Skudlarski P, Fulbright RK, Bronen RA, Fletcher JM, Shankweiler DP, Katz L. Sex differences in the functional organization of the brain for language. *Nature* 1995; 373: 607-609.
- Sullivan EV, Rosenbloom MJ, Desmond JE, Pfefferbaum A. Sex differences in corpus callosum size: relationship to age and intracranial size. *Neurobiol Aging* 2001; 22: 603-611.
- Tang Y, Nyengaard JR, Pakkenberg B, Gundersen HJ. Age-induced white matter changes in the human brain: a stereological investigation. *Neurobiol Aging* 1997; 18: 609-615.
- Tang Y, Whitman GT, Lopez I, Baloh RW. Brain volume changes on longitudinal magnetic resonance imaging in normal older people. *J Neuroimaging* 2001; 11: 393-400.
- Unal B, Kara A, Aksak S, Unal D. A Stereological assessment method for estimating the surface area of cycloids. *EAJM* 2010a; 42: 066-073.
- Unal D, Aksak S, Kara A, Unal B. Östrojen ve Hipokampus İlişkisi. *Turkiye Klinikleri J Neur* 2010b; 5: 167-171.
- Watkins KE, Paus T, Lerch JP, Zijdenbos A, Collins DL, Neelin P, Taylor J, Worsley KJ, Evans AC. Structural asymmetries in the human brain: a voxel-based statistical analysis of 142 MRI Scans. *Cereb Cortex* 2001; 11: 868-877.
- Xu J, Kobayashi S, Yamaguchi S, Iijima K, Okada K, Yamashita K. Gender effects on age-related changes in brain structure. *AJNR Am J Neuroradiol* 2000; 21: 112-118.

