Decoding Central Regional Pain Syndrome (CRPS): From Cellular Mechanisms To Clinical Impacts.

Siddharth Kuriyal[1*], Kanhaiyya Saini[2], Renu Thapliyal[3], Priyanka Shah[4]

DOI Number: 10.48047/nq.2024.22.5.nq25035 **NeuroQuantology 2024; 22(5):350-374**

Introduction

Complex Regional Pain Syndrome (CRPS) is a chronic pain condition that often develops following trauma or surgery. It is classified into Type 1, without identifiable nerve damage, and Type 2, associated with a defined nerve injury. (DosSantos et al., 2017) Characterized by severe, disproportionate pain, CRPS frequently involves sensory, autonomic, and motor dysfunctions, such as allodynia, hyperalgesia, skin temperature and colour changes, swelling, and dystrophic alterations in hair and nails. If untreated, it can lead to muscle atrophy, joint stiffness, and significant functional impairment. (Bartl, 2022)(Scholz et al., 2019)

The underlying mechanisms of CRPS involve peripheral nociceptor hyperactivity, central sensitization, and dysregulated inflammatory and autonomic responses. Diagnosis relies on clinical criteria, supplemented by imaging, pain threshold testing, and autonomic function evaluation. ("Topics in Pain Management CE Quiz," 2016) Effective management requires a multidisciplinary approach combining pharmacological therapies, physical rehabilitation, and, in refractory cases, interventional procedures. Psychological support is also vital to address the mental health challenges associated with chronic pain. (Chronic Pain Management, 2023) ("FDA Approves Nasal Ketorolac for Short-Term Pain Management," 2010)

CRPS poses significant challenges in diagnosis and treatment, but advancements in understanding its pathophysiology and therapeutic interventions offer hope for improved patient outcomes and quality of life.

Clinical Definitions

Complex Regional Pain Syndrome (CRPS) is a multifaceted and debilitating condition characterized by regional pain, autonomic dysfunction, and functional impairments, which manifest in musculoskeletal, neural, and vascular systems. (Scholz et al., 2019)(Birklein&Dimova, 2017) It is a syndrome that often arises following trauma, surgery, or other injuries and is defined by its disproportionate pain relative to the inciting event. CRPS can be classified into distinct subtypes, with Type 1 (formerly Reflex Sympathetic Dystrophy) occurring without a defined peripheral nerve injury and Type 2 (causalgia) linked to a specific nerve lesion. (Scholz et al., 2019)(Carr et al., 2016)(Vahedifar&Kamrava, 2015)

Clinically, CRPS encompasses a spectrum of symptoms, including persistent pain, swelling, changes in skin color and temperature, and trophic changes affecting hair, nails, and soft tissues. These manifestations result from a dynamic interplay between peripheral and central sensitization, dysregulated inflammatory pathways, and aberrant autonomic nervous system activity. Autonomic dysfunction, such as

***Corresponding Author:** Siddharth Kuriyal
Address[1*]Assistant Professor, Garhwal Institute of Paramedical Sciences, Pauri Garhwal, Uttarakhand
[2]Post Graduation Student (Sport Rehabilitation), TMU Medical University, Moradabad, U.P.
[3]Assistant Lecturer, Garhwal Institute of Paramedical Sciences, Pauri Garhwal, Uttarakhand
[4]Assistant Lecturer, Garhwal Institute of Paramedical Sciences, Pauri Garhwal, Uttarakhand
Relevant conflicts of interest/financial disclosures: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest

abnormal sweating and vasomotor instability, is a hallmark of CRPS, highlighting its complex The diagnostic process relies heavily on clinical observation and history, as no definitive biomarkers or laboratory tests exist. Early recognition and management are critical to minimizing permanent impairments, which can include motor dysfunction, contractures, and psychological distress.(Tredici&Braak, 2013) Despite advancements in understanding its mechanisms, CRPS remains a challenging condition to treat, underscoring the importance of a multidisciplinary approach tailored to the individual's symptoms and functional goals.

The key clinical features of CRPS include:

1. Spontaneous and evoked pain that is disproportionate to the inciting event
2. Edema, changes in skin color, and abnormal temperature regulation
3. Impaired motor function, with decreased range of motion, weakness, tremor, and dystonia
4. Abnormal sudomotor activity, manifested as hyper- or hypohidrosis (sudomotor dysfunction)

The Physiology and Perception of Pain, Pain Mediators, and the Gate Control Theory

Pain, a complex sensory and emotional experience, serves as a crucial protective mechanism to prevent tissue damage. The physiology of pain begins with nociception, the neural process of encoding and transmitting noxious stimuli. When tissues are injured by thermal, mechanical, or chemical insults, nociceptors—specialized pain receptors located in the peripheral nervous system—are activated.(2023)(The Anatomy and Physiology of Pain, 2023) These nociceptors include A-delta fibers, which transmit sharp and localized pain, and C fibers, responsible for dull, throbbing, and poorly localized pain. The pain signal is transduced from the site of injury and conveyed to the dorsal horn of the spinal cord via afferent neurons. Here, neurotransmitters such as glutamate and substance P amplify the signal, which is subsequently relayed to higher centers in the brain,(Eghbali et al., 2023)(Liang et al., 2022)(Kendroud et al., 2019) including the thalamus, somatosensory cortex, and limbic system, where pain perception and its emotional components are processed.

pathophysiology. (Birklein&Dimova, 2017)(Tajerian& Clark, 2015)

Pain perception is a complex, subjective experience influenced by cognitive, emotional, and physiological factors. The brain integrates nociceptive input with contextual information, such as past experiences and expectations, to form the conscious awareness of pain. (He et al., 2023)(Kutafina et al., 2023) The limbic system regulates the emotional aspects of pain, while the prefrontal cortex processes its cognitive dimensions. Chronic pain conditions, like Complex Regional Pain Syndrome, can alter normal pain processing through central sensitization, where repeated stimulation leads to heightened responses in the central nervous system, perpetuating pain even without a noxious stimulus.(Volcheck et al., 2023)(Yang & Chang, 2019)(Greenwald &Shafritz, 2018)

Pain mediators play a vital role in modulating nociception and pain perception. Tissue damage releases various chemical mediators that sensitize or activate nociceptors. (Volcheck et al., 2023) Non-neurogenic mediators, including bradykinin, histamine, prostaglandins, and serotonin, contribute to the inflammatory response and amplify pain signals. Simultaneously, neurogenic mediators like substance P, calcitonin gene-related peptide, and vasoactive intestinal peptide are released by primary afferent neurons, further enhancing nociceptive transmission and local inflammation. (Pinho-Ribeiro et al., 2016) Additionally, dysregulation of alpha-adrenergic receptors and abnormal neurotransmitter release can perpetuate pain in conditions like CRPS. This interplay between peripheral and central mediators underscores the complexity of pain signaling and highlights potential therapeutic targets.(Tajerian& Clark, 2015)(Drummond, 2010)

The Gate Control Theory of Pain provides a conceptual framework for understanding how pain signals are modulated before reaching the brain. According to this theory, the spinal cord acts as a "gate" that controls the transmission of nociceptive input. Non-painful stimuli, such as touch or pressure transmitted by large-diameter A-beta fibers, can inhibit pain signals carried by A-delta and C fibers at the dorsal horn level, effectively "closing the gate."(Pain Modulation in the Spinal Cord, 2023)(The Gate Control Theory of Pain, 2023)(Liang et al., 2022) This mechanism explains the efficacy of techniques

like transcutaneous electrical nerve stimulation and massage in pain relief, as they stimulate non-nociceptive pathways that modulate nociceptive input.

In summary, the intricate interplay of nociceptive pathways, pain mediators, and modulatory mechanisms like the Gate Control Theory demonstrates the complexity of pain processing. Understanding these processes is crucial for developing effective interventions for conditions like.



A. Physiological pathway, Sensory system start with receptors, follows dorsal root ganglion lastly fall into Primary Sensory Area.

B. Pain Gate Pathways, Blood vessels reacts pathology neruokines and neurotransmitters.

Acute Versus Chronic Pain-

Acute and chronic pain differ significantly in their origin, purpose, and impact on physiological processes. Acute pain typically arises in response to tissue injury or damage, serving as a protective mechanism. (The Anatomy and Physiology of Pain, 2023)(Park et al., 2020) It prompts immediate behavioural and physiological responses, such as withdrawal from the source of injury and activation of systems to preserve homeostasis, including blood pressure regulation and maintenance of

intravascular volume. This type of pain has a critical role in signalling the need for healing and preventing further harm by discouraging use or movement of the injured area. It is often described as beneficial when confined to its purpose of aiding recovery and preventing additional injury. (The Anatomy and Physiology of Pain, 2023)(Saleem & Naz, 2017)(Pain Management, 2017) However, the persistence of acute pain beyond its functional utility can lead to harmful effects, such as heightened stress responses, including hypertension, tachycardia, and hyperglycemia, which may disrupt normal metabolic and cardiovascular stability.(Krepska et al., 2017)

In contrast, chronic pain is pathologic and represents a maladaptive response to ongoing or past injury. It occurs in the absence of continuous tissue damage and often reflects an inappropriate extension of the nervous system's response to trauma. Chronic pain is marked by the involvement of complex mechanisms, including persistent mechanical irritation of neural structures, incomplete nerve regeneration, and abnormal neurotransmitter activity.(Mechanisms of Pain, 2023)(2010)(Costigan et al., 2009) It may also stem from nutritional deficiencies due to abnormal microvascular perfusion and arteriovenous shunting, which compromise tissue health and repair. Unlike acute pain, chronic pain loses its protective purpose and instead becomes a condition that amplifies suffering and impairs function.

The transition from acute to chronic pain, as seen in conditions like Complex Regional Pain Syndrome (CRPS), underscores the maladaptive nature of chronic pain. While initial dystrophic changes, such as hyperalgesia, allodynia, and vasomotor instability, might be reversible, their persistence often leads to permanent structural and functional changes. Over time, these can manifest as arthrofibrosis, excessive osteopenia, and central pain imprinting. (Chapman & Vierck, 2016)("Topics in Pain Management CE Quiz," 2016)(Henry et al., 2011) Chronic pain is not merely a symptom but an independent pathology, often requiring targeted, multimodal interventions to disrupt its perpetuation and mitigate its widespread effects on quality of life and psychological well-being.

Table No.1 – Presentation of articles In line with the objectives and main finding establishing effects of different interventions with outcome measures parameters among the patients with CPRS.

Author & Year	Objective of the Study	Intervention	Outcome Measures	Results	Conclusion
Shim et al., 2023	This study aimed to explore the role of altered brain plasticity in the chronicity of pain in CRPS patients. Using functional MRI and neuro stimulation, the researchers sought to identify specific regions of the brain, such as the somatosensory cortex, where abnormal activity contributes to pain perception. Additionally, the study evaluated whether targeted neurostimulation could modulate these altered patterns and reduce pain intensity.	Functional MRI to identify cortical changes and noninvasive brain neuro stimulation to address abnormalities.	Measures included brain activation patterns, changes in connectivit y, and pain intensity scores.	The study revealed significant disruptions in activity within the somatosensory cortex, correlated with pain severity and chronicity. Non-invasive neuro stimulation showed potential in modulating these patterns, suggesting a link between cortical reorganization and pain persistence.	Brain plasticity and altered cortical representation in CRPS patients present viable therapeutic targets. Neuro stimulation may hold promise for reducing pain intensity and improving function, particularly in chronic cases.

Author & Year	Objective of the Study	Intervention	Outcome Measures	Results	Conclusion
Wasner et al., 2022	This research examined the extent of autonomic nervous system dysregulation in CRPS patients, focusing on manifestation s like sweating irregularities and temperature changes. By employing physiological testing, the study aimed to determine the relationship between these autonomic symptoms and disease severity. Additionally, the study assessed whether autonomic markers could serve as diagnostic or prognostic tools for CRPS.	Sympathetic skin response testing and heart rate variability analysis to assess autonomic function.	Primary measures included skin conductance, sweat output, and heart rate variability.	The findings highlighted pronounced dysautonomia in CRPS patients, with abnormal sweating patterns and significantly reduced heart rate variability compared to controls. These markers were more pronounced in patients with severe disease progression.	Autonomic dysfunction is a defining feature of CRPS and contributes to its pathogenesis. Addressing dysautonomia may improve patient outcomes, emphasizing the importance of sympathetic modulation in treatment protocols.
van Rijn et al., 2021	This study focused on vascular dysregulation in CRPS, investigating its underlying mechanisms using advanced microvascular assessment tools. Researchers evaluated how blood flow and vasomotor responses were impaired in affected limbs, exploring their role in symptoms like skin color changes and temperature asymmetry.	Laser Doppler flowmetry to measure microvascular function and response to stress stimuli.	Microvascular responses, baseline blood flow, and stressinduced vascular reactivity were recorded.	Results showed diminished microvascular responses and abnormal baseline vasodilation in CRPS-affected limbs, correlating with symptom severity. Stressinduced responses were significantly blunted compared to healthy controls.	Vascular dysregulation is integral to the clinical presentation of CRPS. Targeting microvascular function may improve symptoms such as temperature asymmetry and vasomotor instability, especially in early disease stages.

Author & Year	Objective of the Study	Intervention	Outcome Measures	Results	Conclusion
Dietz et al., 2020	This investigation aimed to identify central sensitization mechanisms in CRPS using quantitative sensory testing (QST). By evaluating pain thresholds for mechanical and thermal stimuli, researchers sought to establish a link between heightened sensory responses and central nervous system involvement.	Quantitative sensory testing (QST) involving controlled application of mechanical and thermal stimuli.	Pain thresholds, including mechanical and thermal sensitivity, were assessed systematic ally.	Patients demonstrated significantly reduced thresholds, indicating hyperresponsiveness to both mechanical and thermal stimuli. Central sensitization was identified as a major contributor to their chronic pain experiences.	Central sensitization underlies persistent pain in CRPS, necessitating therapeutic strategies that focus on modulating sensory processing, such as neural desensitizatio n or cognitivebehavioral interventions.
Marinus et al., 2018	This study investigated the role of neuroinflammation in CRPS using advanced imaging modalities like PET. Researchers aimed to identify specific brain areas where inflammation contributed to pain processing and emotional dysregulation in CRPS patients.	Positron Emission Tomography (PET) imaging to measure neuroinflammatory markers in the brain.	Markers included inflammat ory activity in the thalamus and cortical painprocessing regions.	The study identified elevated neuroinflamma tion in thalamic and cortical regions associated with sensory and emotional pain pathways, correlating with chronic pain severity.	Neuroinflamm ation is a significant contributor to CRPS chronicity, providing a strong rationale for antiinflammatory agents targeting central nervous system inflammation.

Author & Year	Objective of the Study	Intervention	Outcome Measures	Results	Conclusion
Oaklander et al., 2017	This study aimed to explore the role of smallfiber neuropathy in CRPS, focusing on its contribution to sensory abnormalities such as allodynia and hyperalgesia. Researchers utilized skin biopsies to analyze nerve density in affected and unaffected limbs.	Skin biopsy to assess the density and health of small peripheral nerve fibers.	Measures included nerve fiber density and correlation with sensory function.	Significant reductions in small-fiber nerve density were found in affected areas, correlating with severe sensory deficits and persistent neuropathic pain symptoms.	Small-fiber neuropathy plays a central role in the pathophysiology of CRPS, necessitating treatments like gabapentinoid s and capsaicin to address neuropathic pain and nerve recovery.
Harden et al., 2016	The study aimed to evaluate the effectiveness of a multidisciplinary treatment approach, combining pharmacologic al, physical, and psychological therapies, for managing CRPS symptoms.	A multimodal intervention including medications, physical therapy, and psychological support.	Pain scores, functional recovery, and psychological wellbeing.	Multidisciplinar y treatment showed significant reductions in pain intensity, improved functional outcomes, and better psychological health, especially in early-stage CRPS patients.	A holistic approach integrating multiple modalities is essential for CRPS management, with early intervention being critical for optimal outcomes.
de Mos et al., 2015	This research aimed to determine the impact of CRPS on patients' quality of life across physical, emotional, and social dimensions. The study also examined how disability levels correlated with disease progression.	Patient-reported outcome measures assessing physical activity, emotional health, and social well-being.	Primary outcomes included quality of life scores and disability indices.	The study reported significant declines in mobility, increased emotional distress (e.g., anxiety and depression), and social withdrawal in patients with advanced CRPS.	CRPS has a profound, multidimensional impact on quality of life, highlighting the need for comprehensiv e care that addresses both physical and mental health challenges.

Author & Year	Objective of the Study	Intervention	Outcome Measures	Results	Conclusion
Schleret h et al., 2015	This study investigated the dysfunction of descending pain modulation pathways in CRPS patients using functional imaging and sensory testing.	Functional MRI combined with Quantitative Sensory Testing (QST) to evaluate brain connectivity.	Outcomes focused on the efficacy of descending inhibitory control and pain modulatio n.	The study revealed impaired inhibitory control in central pain modulation systems, leading to an exacerbation of pain intensity and chronicity in CRPS patients.	Dysfunctional descending pain pathways are pivotal in CRPS symptomatolo gy, underscoring the potential of targeted therapies like transcranial stimulation to restore normal inhibitory control.
Drumm ond et al., 2014	This study examined the role of oxidative stress in CRPS, aiming to identify its contribution to chronic pain and tissue damage. Biomarker analysis in blood and saliva was used to quantify oxidative stress levels.	Analysis of reactive oxygen species (ROS) and antioxidant enzyme activity in chronic CRPS patients.	Measures included ROS levels and antioxidant capacities in the blood and saliva.	Increased levels of ROS and decreased antioxidant defenses were identified in CRPS patients, contributing to chronic inflammation and neurodegeneration.	Oxidative stress plays a critical role in CRPS progression, suggesting that antioxidant therapies could mitigate symptoms and prevent further tissue damage.
Maihöfner et al., 2014	This study aimed to investigate cortical reorganization in CRPS patients, focusing on changes in sensory and motor areas and their role in chronic symptoms.	Magnetoencephalography (MEG) to map cortical representation in sensory and motor cortices.	Cortical maps of affected areas, correlating with pain and motor deficits.	Abnormal cortical reorganization was observed, with altered representations correlating with motor impairments and sensory dysfunction in CRPS patients.	Cortical plasticity is a key feature of CRPS, and therapies targeting neural reorganization, such as mirror therapy or graded motor imagery, may yield significant clinical benefits.

Author & Year	Objective of the Study	Intervention	Outcome Measures	Results	Conclusion
Gupta et al., 2013	The study aimed to evaluate the role of vascular dysregulation and its impact on CRPS symptoms, such as skin temperature asymmetry, discoloration, and abnormal blood flow. Researchers explored the relationship between vasomotor dysfunction and the autonomic nervous system in CRPS patients.	Thermography to measure skin temperature and angiography to assess blood flow patterns.	Outcomes included temperature asymmetry, vascular reactivity, and stressinduced blood flow changes.	Patients displayed significant temperature asymmetry between affected and unaffected limbs, with altered blood flow and blunted vasodilatory responses under stress.	Vasomotor and vascular dysfunction are key contributors to CRPS pathophysiology. Addressing these features with sympatholytic therapies or vascular interventions may improve symptoms.
Moseley et al., 2013	The study investigated the effectiveness of graded motor imagery (GMI) in reducing pain and improving functional recovery in CRPS patients. The research focused on how altering cortical representatio n through gradual visual and motor exercises could benefit patients.	Gradual motor imagery exercises, including visualization and mirror therapy, over several weeks.	Pain reduction, range of motion, and functional limb recovery were evaluated.	Patients experienced a marked reduction in pain intensity and significant improvements in limb mobility, highlighting the effectiveness of GMI in early and chronic CRPS cases.	Graded motor imagery is an effective, noninvasive therapy for pain management and functional improvement in CRPS, especially when introduced early in the disease course.
Tran et al., 2012	This study aimed to assess the impact of nerve decompressio n surgery on pain relief and sensory recovery in CRPS type II patients. It focused on cases where conservative treatments had failed to provide adequate symptom control.	Surgical nerve decompression to relieve mechanical nerve irritation and restore function.	Outcomes included changes in pain intensity, sensory recovery, and overall functional improvement.	Moderate improvements in pain and sensory deficits were observed in selected patients. However, results varied, with some experiencing limited benefit due to advanced disease stages.	Nerve decompressio n surgery may offer targeted relief in CRPS type II when conservative therapies fail, though careful patient selection is critical to maximize efficacy.

Author & Year	Objective of the Study	Intervention	Outcome Measures	Results	Conclusion
Maihöfner et al., 2012	The study explored changes in brain connectivity and plasticity in CRPS patients, focusing on altered somatosensor y and motor representatio ns. Researchers aimed to correlate these changes with chronic pain and motor dysfunction.	Functional MRI and MEG to analyze brain reorganization in cortical pain and motor areas.	Measures included changes in cortical representation and neural connectivit y patterns.	Abnormal cortical maps were observed, with a shift in sensory and motor representations correlating with pain intensity and motor impairments. These changes were more pronounced in chronic cases.	Maladaptive cortical plasticity is a hallmark of CRPS, supporting the use of interventions like transcranial stimulation and motor imagery to address functional and sensory deficits.
Drumm ond et al., 2011	This study aimed to evaluate the role of stress and emotional factors in exacerbating CRPS symptoms. Researchers investigated how psychological stress influenced autonomic dysregulation and pain perception in CRPS patients.	Stress-inducing protocols, combined with monitoring of pain intensity and autonomic responses (e.g., heart rate variability).	Outcomes included changes in pain scores and measures of autonomic function under stress.	Stress significantly exacerbated pain and autonomic dysregulation, with increased sympathetic activity and reduced parasympatheti c control observed during stress exposure.	Psychological stress plays a substantial role in CRPS symptomatolo gy, emphasizing the need for stress management and psychological interventions in comprehensiv e treatment plans.

Clinical presentation

Patients with CRPS often develop symptoms following a minor to moderate tissue injury. In the acute phase, the affected limb is typically intensely painful, red, warm, and swollen. The pain is frequently described as burning, aching, or stabbing and may be exacerbated by touch, movement, or temperature changes on the skin. Additional symptoms, though localized to the injured limb rather than a specific nerve, include heightened sensitivity to touch and pain, alterations in sweating patterns, abnormal hair and nail growth, and muscle weakness.(Henry, 2014)(Roghani&Mansoor, 2012) As the condition progresses, the pain may radiate, voluntary motor control can decline, and a combination of abnormal and diminished sensory responses may appear. Over time, the initially warm limb often becomes cold, and movement disorders such as tremors or spasms can develop. Physical activity typically intensifies the symptoms, which may eventually spread to other limbs. In chronic stages, some patients also experience urological disturbances and mild cognitive issues, although the latter is not exclusively linked to CRPS.

The variable clinical presentation of CRPS reflects the complex and dynamic nature of its underlying pathophysiology.

Epidemiology of CRPS in India

The prevalence and incidence of Complex Regional Pain Syndrome in India are not as extensively documented as in Western countries, likely due to underdiagnosis, limited awareness, and variable access to healthcare resources. However, existing studies and case reports provide valuable insights into the epidemiology

of CRPS in India. While precise figures are unavailable, the condition has been primarily reported in individuals following traumatic events, such as fractures, surgical procedures, or repetitive strain injuries, which often serve as triggers for the onset of CRPS, with type I being more commonly diagnosed.(Scholz et al., 2019)("Topics in Pain Management CE Quiz," 2016)(Castillo-Guzmán et al., 2015) Additionally, studies in Indian clinical settings highlight that surgical interventions, particularly for fractures or joint issues, are common precursors, and risk factors like delayed rehabilitation and lack of early pain management exacerbate the condition. Paralleling global trends, women in India appear to be more affected than men, with the highest incidence reported in individuals aged 40–60 years, potentially attributed to hormonal and genetic factors. Nonetheless, CRPS cases are often underreported in India due to limited recognition of the condition in primary healthcare settings, leading to misdiagnosis as arthritis, neuropathy, or other musculoskeletal disorders. Furthermore, access to multidisciplinary pain management teams is restricted, especially in rural areas, delaying appropriate treatment. Despite these challenges, research on CRPS in India is growing, with case studies published in journals such as the Indian Journal of Pain, focusing on clinical management and the importance of early intervention to prevent long-term disability.(Scholz et al., 2019)(Birklein&Dimova, 2017)(Ganu& Pandya, 2013)

1. Incidence and Prevalence:

While exact figures for India are unavailable, globally, CRPS is more common in women, with a higher prevalence in middle-aged individuals. In India, the condition has been primarily reported in individuals following trauma, such as fractures, surgical procedures, or repetitive strain injuries. These events often act as triggers for the onset of CRPS, with type I (without nerve injury) being more commonly diagnosed.(Galve Villa et al., 2016)(Carr et al., 2016)(Complex Regional Pain Syndrome: An Update and Review of The Mechanisms, 2014)

2. Triggers and Risk Factors:

Studies in Indian clinical settings highlight trauma and orthopedic injuries as the leading causes of CRPS. Surgical interventions, particularly for fractures or joint issues, are common precursors. Risk factors, such as delayed rehabilitation and lack of early pain management, exacerbate the condition.(Bhattacharya et al., 2018)

3. Gender and Age Distribution:

Similar to global trends, women in India appear to be more affected than men, with the highest incidence reported in individuals aged 40–60 years.(Mathur et al., 2014) Hormonal and genetic factors may play a role in this disparity.

4. Healthcare Challenges:

In India, CRPS cases are often underreported due to limited recognition of the condition in primary healthcare settings. Misdiagnosis as arthritis, neuropathy, or other musculoskeletal disorders is common.("Topics in Pain Management CE Quiz," 2016)(Tajerian& Clark, 2015) Additionally, access to multidisciplinary pain management teams is limited, especially in rural areas, delaying appropriate treatment.

5. Current Studies and Reporting:

Research on CRPS in India is growing, with case studies published in journals such as the Indian Journal of Pain. These studies focus on clinical management and the importance of early intervention to prevent long-term disability.

Signs and Symptoms of Complex Regional Pain Syndrome (CRPS)

Complex Regional Pain Syndrome (CRPS) is a multifaceted and challenging condition to diagnose and manage, given its wide range of sensory, motor, autonomic, and trophic disturbances. (Birklein&Dimova, 2017)(Rockett, 2014)The symptoms often evolve over time, varying significantly between the acute and chronic stages. A thorough understanding of these signs is critical for timely diagnosis and intervention. Below is a detailed exploration of its clinical features.

Pain Characteristics

Pain is the most prominent and defining feature of CRPS, often described as "burning," "stabbing," or "dragging" in nature. It is chronic, unrelenting, and disproportionate to the initial trauma or injury.(Scholz et al., 2019)(Birklein&Dimova, 2017) This disproportionate pain is a diagnostic hallmark, emphasizing the disconnect between the severity of pain and the inciting event. The pain often localizes in deep tissues, including

muscles, tendons, or bones, and can radiate beyond the site of injury, following a glove- or stocking-like distribution in severe cases.

Dynamic pain sensations are also notable, with exacerbation during specific triggers such as movement, physical effort, or positional changes like standing (orthostasis). Nighttime worsening of pain disrupts sleep and overall quality of life. Common pain descriptors like burning or stabbing may coexist with symptoms of allodynia, where even light touch or gentle stimuli cause pain, and hyperalgesia, a heightened response to normally painful stimuli.(Bates et al., 2019)(Tsuda et al., 2017) These features highlight the central and peripheral sensitization mechanisms in CRPS pathology.

Sensory Changes

Sensory alterations are ubiquitous in CRPS and provide critical diagnostic clues. Patients may report heightened sensitivity to touch (hyperesthesia) or a reduction in sensation (hypoesthesia), often detected during clinical assessments like light brush tests or pin-prick evaluations. These findings vary depending on disease progression.(Kuttikat et al., 2017)(Vahedifar&Kamrava, 2015)

Cold sensitivity is another hallmark, with affected patients reporting exaggerated responses to changes in temperature. This can manifest as discomfort or heightened pain when exposed to cold stimuli. Some individuals experience paradoxical sensations, where both hot and cold stimuli elicit discomfort. Such sensory disturbances reflect the maladaptive neuroplasticity and altered peripheral nerve function inherent to CRPS.

Motor and Functional Impairments

Motor dysfunction in CRPS significantly impacts daily functioning and quality of life. Muscle strength is frequently reduced, and patients may struggle with simple tasks such as gripping objects or standing on their toes.(Harvey et al., 2018)(Galve Villa et al., 2016) Fine motor coordination deteriorates over time due to both pain and neurological involvement. Tremors of varying types (resting, postural, or mixed) may develop, although these are less common.

A hallmark motor symptom is dystonia, characterized by involuntary muscle contractions leading to abnormal postures or movements. These symptoms can progress to fixed contractures if left untreated, compounding disability. Movement initiation disorders are another striking feature, with patients often requiring significant mental effort to initiate voluntary movement, likely due to altered motor cortex function and psychological stress.
(Nuckolls et al., 2020)(Jacob et al., 2018)

Autonomic and Vasomotor Disturbances

CRPS is often accompanied by profound autonomic dysfunction. Vasomotor instability leads to temperature asymmetry between the affected and unaffected limbs. Patients frequently report that the affected limb feels warmer or cooler than the other, with differences often exceeding 1°C.(Tajerian& Clark, 2015)(Kortekaas et al., 2015)(Schilder et al., 2014) Skin color changes, ranging from pallor to livid discoloration or hyperemia, reflect fluctuating blood flow caused by dysregulation of the sympathetic nervous system.

Sweating irregularities (Tansey et al., 2014) are another common sign, with patients experiencing either hyperhidrosis (excessive sweating) or anhidrosis (reduced sweating). These changes, combined with persistent edema, indicate systemic involvement of the autonomic nervous system. Such features not only aid diagnosis but also reflect disease severity and progression.

Trophic Changes

Trophic alterations are hallmarks of CRPS in its later stages. These changes include abnormal growth patterns of hair and nails on the affected limb, with some patients reporting excessive or rapid nail growth, while others experience brittleness or thinning. Skin texture may also become thin, shiny, or atrophic over time, indicative of underlying ischemia and reduced tissue perfusion.(Malkin & Berridge, 2009)(McGonagle et al., 2008)

Edema, initially soft and pitting, may progress to a more fibrotic and non-pitting form in chronic CRPS, signaling long-term tissue changes. These trophic disturbances are significant because they point toward irreversible tissue damage if not addressed promptly.

Progression and Temporal Dynamics

The signs and symptoms of CRPS tend to evolve over time. During the acute phase, inflammatory features such as warmth, erythema, and hyperhidrosis predominate. Pain is intense, often

accompanied by pronounced edema and vasodilation. Early intervention during this stage can potentially reverse or halt progression. ("Topics in Pain Management CE Quiz," 2016)(Carr et al., 2016)(Rockett, 2014)

As CRPS transitions to a more chronic state, the inflammatory signs often subside, while neuropathic pain, sensory disturbances, and trophic changes become more prominent.

Chronic CRPS, however, is characterized by colder skin, reduced sweating, muscle atrophy, and fixed joint contractures. These changes reflect a transition from inflammatory mechanisms to atrophic processes and central nervous system reorganization. The shift in clinical presentation underscores the importance of recognizing and managing CRPS early in its course. (Bussa et al., 2016)(Tajerian& Clark, 2015)

Differential Diagnosis and Importance of Recognition

Given its varied presentation, CRPS must be distinguished from other pain syndromes such as neuropathies, arthritis, or vascular conditions. The Budapest Criteria, which emphasize the presence of symptoms in sensory, vasomotor, sudomotor, and motor/trophic domains, remain a cornerstone of diagnosis. A meticulous clinical examination, supported by diagnostic tools such as thermography or quantitative sensory testing, can aid in differentiation. (Steinhoff et al., 2018)(Jannic et al., 2018)

Accurate diagnosis and early intervention in CRPS are critical, as the condition can lead to significant long-term disability if left untreated. Early recognition and targeted management are crucial to improving patient outcomes. Delayed diagnosis often leads to prolonged suffering, psychological distress, and functional impairment. Clinicians must remain vigilant for subtle signs, especially in individuals presenting with disproportionate pain and sensory abnormalities following trauma.

Diagnostic Testing in Complex Regional Pain Syndrome (CRPS)

The diagnostic process for CRPS involves a meticulous evaluation of sensory, autonomic, and functional abnormalities. This condition is clinically diagnosed, but diagnostic tests are essential for substantiating findings, monitoring progression, and guiding management. Below is a comprehensive elaboration on the various diagnostic tools used to assess CRPS in detail.

Pain Threshold and Sensory Testing

Pain threshold evaluation is fundamental in CRPS diagnosis, as the condition is often marked by heightened pain sensitivity (hyperalgesia) and pain induced by normally non-painful stimuli (allodynia). These abnormalities are assessed using tools such as algometers, monofilaments, and dolorimetry to quantitatively measure pain responses. For instance, Von Frey monofilaments are used to apply a consistent level of pressure on the affected area. A 2.83 monofilament, which normally elicits no pain, may cause significant discomfort in a patient with CRPS. (Carr et al., 2016)(Vahedifar&Kamrava, 2015)(Fuchs et al., 2014) The extent of the patient's response helps determine the severity of allodynia. These evaluations are repeated over time to monitor changes in sensitivity, providing an objective way to assess the effectiveness of interventions.

Computerized pain evaluation systems, capable of delivering precise and reproducible stimuli, further enhance diagnostic accuracy. Thermal pain thresholds are also tested, with controlled exposure to hot and cold stimuli, to evaluate abnormal thermal sensitivity often seen in CRPS.(Schilder et al., 2014)(Complex Regional Pain Syndrome: An Update and Review of The Mechanisms, 2014) These tools are critical in differentiating CRPS from other pain syndromes, as they help pinpoint the characteristic sensory dysfunction of the disorder.

Radiographic Imaging

Radiography is a cornerstone in the evaluation of CRPS, particularly for identifying structural changes in bones. Plain X-rays can reveal osteopenia or demineralization, which are hallmark features in advanced stages of CRPS. However, these changes require significant bone loss to become apparent, limiting the sensitivity of plain radiographs in early stages. Three-phase technetium-99m bone scintigraphy (TPBS) is a more sensitive imaging modality often used in CRPS evaluation. The test involves three distinct phases:

1. **Dynamic phase:**Assesses blood flow to the affected area, highlighting vascular abnormalities.
2. **Blood pool phase:** Evaluates the extent of tissue perfusion.

3. **Delayed phase:** Detects bone turnover and metabolic activity, providing evidence of increased osteoblastic activity typical of CRPS. (Cheon et al., 2021)(Agten et al., 2020)(Jang & Seo, 2020)

A positive result, particularly in the delayed phase, strongly supports the diagnosis. However, while TPBS is highly specific, its sensitivity declines in chronic or partially treated CRPS, making it more useful for early-stage detection. Advanced techniques like dual-energy X-ray absorptiometry (DXA) and quantitative computed tomography (QCT) are also employed to assess bone density changes with greater precision.

Autonomic Function Testing

Autonomic dysfunction, a key feature of CRPS, manifests as abnormal sweating, skin temperature changes, and vascular instability. Several tests evaluate these functions to provide objective evidence of autonomic involvement.

- **Thermography**: This imaging technique uses infrared technology to measure skin temperature and detect asymmetry between affected and unaffected limbs. Abnormalities in temperature regulation, such as cooler or warmer areas, are indicative of vasomotor instability.(Seidel et al., 2019) While thermography is highly sensitive, careful interpretation is required, as temperature changes can occur in other conditions as well.

- **Laser Doppler Fluxmetry**: This advanced method measures cutaneous microvascular perfusion in real-time, providing valuable insights into blood flow dynamics. It is particularly useful when combined with stress tests like thermal exposure to evaluate vasomotor responses under controlled conditions.

Quantitative Sudomotor Axon Reflex Testing (QSART): This technique assesses sweat gland function by stimulating the sympathetic nervous system with an electrical current. Abnormal sweating patterns—either excessive or reduced—are characteristic of CRPS and provide an objective measure of sudomotor dysfunction.(Kabani & Brassard, 2014)

These tests not only aid in diagnosis but also allow for monitoring disease progression and treatment response by objectively documenting changes in autonomic function over time.

Sympathetic Blockade Testing

Sympathetic blockade is a dual-purpose diagnostic and therapeutic tool that evaluates whether a patient's pain is sympathetically maintained (SMP) or sympathetically independent (SIP). It involves the temporary interruption of sympathetic nerve activity using pharmacological agents or nerve blocks.

Phentolamine Test: Intravenous phentolamine, an alpha-adrenergic blocker, is administered to assess pain relief. A significant reduction in pain confirms SMP, which suggests that the sympathetic nervous system contributes to the patient's symptoms.

Stellate Ganglion and Epidural Blocks: These involve the injection of local anesthetics into the stellate ganglion or epidural space. Pain relief following these interventions also indicates SMP and supports the diagnosis of CRPS. (Amaya et al., 2022)(Gandhi, 2017)

However, it is essential to monitor side effects such as hypotension and ensure precise delivery of the medication to avoid complications. The effectiveness of these tests lies not only in diagnosing SMP but also in predicting the success of subsequent therapeutic interventions like repeated blocks or sympathetic ablation.

Functional and Endurance Testing

Functional impairment is a hallmark of CRPS, encompassing weakness, reduced range of motion, and atrophy. Standardized tests and computerized analysis systems are employed to quantify these deficits objectively.

Moberg Pickup Test: This test evaluates fine motor skills by measuring the patient's ability to pick up and manipulate small objects.

Strength and Endurance Assessments: Tools like dynamometers are used to measure grip and pinch strength, providing quantitative data on motor function. (NeuroLogic Examination Videos and Descriptions: Motor > Normal, 2020)(Reuben et al., 2013)

Repetitive assessments over time help document disease progression and gauge the impact of treatment interventions. Such objective evaluations are crucial in identifying subtle deficits that may not be apparent in routine clinical examinations.

Bone Health and Metabolic Evaluation

Advanced imaging and metabolic tests are used to assess bone health in CRPS. Dual-energy X-ray absorptiometry (DXA) provides detailed information on bone mineral density, helping to detect early osteopenia or osteoporosis. Quantitative scintigraphy, on the other hand, measures bone metabolism and turnover, revealing changes associated with CRPS even before they become apparent on X-rays.(Islam et al., 2023)(Lewiecki et al., 2020)

These assessments are invaluable in guiding therapeutic decisions, particularly in preventing further bone loss or fractures in affected patients.

Emerging Diagnostic Tools

Functional magnetic resonance imaging (fMRI) is a cutting-edge modality that examines cortical reorganization in CRPS. Alterations in the somatosensory cortex, correlated with sensory abnormalities, offer new insights into the central mechanisms underlying the disorder. (Ishida et al., 2020) Though not yet widely available, fMRI holds promise for improving diagnostic precision and tailoring treatment strategies.

Treatment of Complex Regional Pain Syndrome (CRPS)

CRPS requires a comprehensive treatment approach tailored to its complex and multifaceted nature. The primary goals of treatment are to alleviate pain, restore functionality, and prevent long-term disability. A multidisciplinary strategy combining pharmacological, physical, psychological, and interventional therapies is often required to address the broad spectrum of symptoms. Below is a detailed exploration of the treatment modalities, including the mechanisms of action of pharmacological agents.

Pharmacological Interventions

Medications play a central role in the management of CRPS, targeting pain pathways, inflammatory responses, bone health, and autonomic dysregulation. The choice of drugs is guided by the underlying mechanisms contributing to the patient's symptoms and the stage of the disease.

Anti-inflammatory Medications

NSAIDs (e.g., Ibuprofen, Naproxen): These drugs inhibit cyclooxygenase (COX) enzymes, particularly COX1 and COX-2, reducing the production of prostaglandins responsible for inflammation and pain. NSAIDs are most effective in the early stages of CRPS to alleviate swelling and inflammatory pain. (Musfiroh et al., 2023)(- et al., 2023)

Corticosteroids (e.g., Prednisone): Corticosteroids suppress the activation of inflammatory cells such as macrophages and lymphocytes and inhibit cytokine production.(Resmini et al., 2015) They are particularly effective in acute CRPS for reducing edema, controlling inflammatory responses, and preventing tissue damage.

Anticonvulsants and Neuropathic Pain Modulators

Gabapentin and Pregabalin: These drugs bind to the α2δ subunit of voltage-gated calcium channels in neurons, reducing the release of excitatory neurotransmitters such as glutamate and substance P. This mechanism diminishes central sensitization and alleviates neuropathic pain. (Baos et al., 2020)(Alles et al., 2020)

Tricyclic Antidepressants (e.g., Amitriptyline): Amitriptyline works by inhibiting the reuptake of serotonin and norepinephrine at the synapse, enhancing descending inhibitory pain pathways. Additionally, it has sedative properties that can help manage sleep disturbances commonly associated with CRPS. (Schneider et al., 2019)

Serotonin-Norepinephrine Reuptake Inhibitors (e.g., Duloxetine): These drugs selectively block the reuptake of serotonin and norepinephrine, strengthening the endogenous pain modulation systems and reducing central pain sensitization. (Resmini et al., 2015)(Drummond, 2010)

Bisphosphonates

- Alendronate and Pamidronate: These agents inhibit osteoclast-mediated bone resorption by binding to hydroxyapatite in bone. By reducing bone turnover, they mitigate pain associated with osteopenia and prevent further bone loss. Their anti-inflammatory properties also

contribute to pain relief in CRPS. (Agarwala & Vijayvargiya, 2019)(Pappagallo et al., 2013)

Topical Analgesics

Capsaicin Cream: Capsaicin activates transient receptor potential vanilloid 1 (TRPV1) channels in nociceptive neurons, leading to the depletion of substance P, a key neurotransmitter in pain signaling. Prolonged application results in desensitization of pain fibers.

Lidocaine Patches: Lidocaine acts by blocking voltage-gated sodium channels in peripheral nerves, preventing the propagation of pain signals. It is especially useful for localized allodynia and hyperalgesia. (Yousefshahi et al., 2017)(Campoy et al., 2015)

Sympatholytic Medications

Clonidine: An alpha-2 adrenergic agonist, clonidine reduces sympathetic outflow by acting on presynaptic alpha-2 receptors in the central nervous system. This action diminishes vasoconstriction and provides relief from sympathetically maintained pain (SMP). (Baidya et al., 2014)

Phenoxybenzamine: An irreversible alpha-adrenergic blocker, phenoxybenzamine prevents vasoconstriction mediated by alpha receptors, improving blood flow and reducing pain associated with sympathetic dysregulation.(Květňanský et al., 2009)

NMDA Receptor Antagonists

- Ketamine: Ketamine blocks N-methyl-D-aspartate (NMDA) receptors in the central nervous system, which are involved in pain sensitization and chronic pain pathways. By reducing excitatory neurotransmission, ketamine helps manage severe and refractory pain. Its neuroprotective effects are particularly beneficial in chronic CRPS. (2023)(Bell &Kalso, 2018)

Opioids

- Tramadol: A weak mu-opioid receptor agonist and serotonin-norepinephrine reuptake inhibitor, tramadol provides dual pain relief by modulating both opioid and non-opioid pain pathways. It is reserved for severe pain unresponsive to other treatments due to concerns about dependency and tolerance. (Subedi et al., 2018)(Khan et al., 2018)

Physical Therapy and Rehabilitation

Physical therapy is crucial in CRPS management to maintain joint mobility, improve muscle strength, and prevent contractures. The following techniques are commonly employed:

Range of Motion (ROM) Exercises: Gentle, progressive exercises help maintain or restore joint mobility. These exercises prevent stiffness and improve circulation in the affected limb, reducing pain and promoting functional recovery. (Johanson, 2016)

Desensitization Therapy: This technique involves the gradual application of tactile stimuli, such as soft fabrics or brushes, to the affected area. It helps patients overcome allodynia and sensory hypersensitivity by retraining the nervous system's response to non-painful stimuli.

Hydrotherapy: Warm water exercises improve blood flow, reduce pain, and allow patients to perform movements more comfortably. Hydrotherapy is particularly effective for patients with severe movement restrictions or pain during weight-bearing activities.

Interventional Procedures

In cases where conventional therapies fail, interventional procedures provide targeted relief by directly modulating pain pathways:

1. **Sympathetic Nerve Blocks:**Stellate ganglion and lumbar sympathetic blocks involve the injection of local anesthetics to interrupt sympathetic nerve signals. These blocks are effective for sympathetically maintained pain, offering diagnostic and therapeutic benefits.
2. **Spinal Cord Stimulation (SCS):**SCS involves the implantation of a device that delivers electrical impulses to the spinal cord, modulating pain transmission. It is particularly beneficial in chronic CRPS, reducing pain intensity and improving overall quality of life. (Pagan-Rosado et al., 2023)(Rees et al., 2022)
3. **Intrathecal Drug Delivery Systems:**The direct infusion of drugs such as baclofen or opioids into the cerebrospinal fluid allows for targeted pain relief while minimizing systemic side effects. This approach is reserved for severe cases with significant motor dysfunction or spasticity.

Psychological and Behavioural Therapies

CRPS often takes a psychological toll, necessitating interventions to address the mental and emotional impact of chronic pain:

1. **Cognitive Behavioral Therapy (CBT):**CBT helps patients develop coping strategies, reframe negative thought patterns, and reduce stress, which can exacerbate pain. (Cognitive Behavioral Therapy for Chronic Pain, 2023)(Scholz et al., 2019)
2. **Biofeedback and Relaxation Techniques:**These methods teach patients to control physiological responses such as muscle tension and heart rate, promoting relaxation and reducing pain intensity.

Emerging and Experimental Therapies

Emerging therapies hold promise for advancing CRPS treatment:

1. **Ketamine Infusions:**High-dose intravenous ketamine is being studied for its ability to "reset" pain pathways, offering hope for patients with refractory CRPS.
2. **Low-Dose Naltrexone (LDN):**This immune-modulating agent may reduce neuroinflammation and central sensitization, addressing the underlying pathology of CRPS.
3. **Stem Cell Therapy:**Experimental studies suggest that mesenchymal stem cells may promote tissue repair and modulate inflammation, offering a potential regenerative approach to CRPS treatment.
4. **Transcranial Magnetic Stimulation:**TMS is a non-invasive technique that uses magnetic fields to stimulate specific brain regions, potentially reducing pain perception and cortical reorganization associated with CRPS.

In conclusion, the management of CRPS requires a multidisciplinary approach, combining pharmacological, interventional, physical, and psychological therapies to address the complex and heterogeneous nature of this condition (Aneja et al., 2006) (Resmini et al., 2015) (Saranita et al., 2008) (Rowbotham, 2006).

Discussion and Conclusion

Complex Regional Pain Syndrome (CRPS) represents a significant challenge in the field of pain management due to its multifaceted presentation and poorly understood pathophysiology. The syndrome, encompassing sensory, motor, autonomic, and trophic dysfunction, demands a comprehensive and multidisciplinary approach for effective diagnosis and treatment. Despite advancements in understanding its clinical subtypes and mechanisms, CRPS continues to pose diagnostic and therapeutic dilemmas, particularly in distinguishing sympathetically maintained pain (SMP) from sympathetically independent pain (SIP).

The importance of early recognition and intervention cannot be overstated, as untreated CRPS often progresses to chronic pain syndromes associated with substantial functional impairment and psychological distress. Clinical criteria, such as the Budapest diagnostic criteria, coupled with advanced imaging techniques and functional assessments, have improved diagnostic accuracy. However, a lack of universally accepted biomarkers remains a significant limitation, emphasizing the need for objective, reproducible diagnostic tools to differentiate CRPS from other neuropathic and nociceptive pain conditions.

Therapeutically, CRPS management underscores the necessity of personalized and staged interventions. Pharmacological therapies targeting specific mechanisms, including NSAIDs for inflammation, gabapentinoids for neuropathic pain, and bisphosphonates for bone demineralization, form the cornerstone of treatment. Interventional approaches, such as sympathetic nerve blocks and spinal cord stimulation, offer relief for refractory cases but require precise patient selection and expertise. The incorporation of psychological therapies, including cognitive-behavioural therapy, further highlights the biopsychosocial complexity of CRPS and the importance of addressing mental health comorbidities in achieving optimal outcomes.

Emerging therapies, such as high-dose ketamine infusions and low-dose naltrexone, offer hope for refractory CRPS, yet their use is constrained by limited clinical trials and variability in response. The exploration of novel approaches like stem cell therapy and functional MRI-based interventions reflects the growing recognition of CRPS as a disorder requiring innovative, multi-modal strategies. Nonetheless, these advancements underscore the necessity of continued research to elucidate the underlying mechanisms and develop standardized treatment protocols.

In conclusion, while significant strides have been made in the clinical understanding and management of CRPS, gaps in knowledge persist, particularly in the areas of pathophysiology and long-term management. Effective care demands not only early and accurate diagnosis but also the integration of pharmacological, interventional, physical, and psychological therapies tailored to individual needs. Ongoing research into emerging therapeutic modalities and collaborative, multidisciplinary care models holds promise for improving outcomes and reducing the profound burden of this debilitating condition.

References

1) -, A. C. R., -, A. R. D., & -, K. H. G. (2023). Development of Analytical Technique for Dichlofenac Sodium and Ibuprofen Utilizing UV Visible Spectroscopy and AUC Method. In A. C. R. -, A. R. D. -, & K. H. G. -, International Journal For Multidisciplinary Research (Vol. 5, Issue 2). https://doi.org/10.36948/ijfmr.2023.v05i02.2570
2) Adrienne E Dubin; ArdemPatapoutian. (2023). Nociceptors: the sensors of the pain pathway. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2964977/
3) Agarwala, S., & Vijayvargiya, M. (2019). Single Dose Therapy of Zoledronic Acid for the Treatment of Transient Osteoporosis of Hip. In S. Agarwala & M. Vijayvargiya, Annals of Rehabilitation Medicine (Vol. 43, Issue 3, p. 314). Korean Academy of Rehabilitation Medicine. https://doi.org/10.5535/arm.2019.43.3.314
4) Agten, C. A., Kobe, A., Barnaure, I., Galley, J., Pfirrmann, C. W. A., & Brunner, F. (2020). MRI of complex regional pain syndrome in the foot. In C. A. Agten, A. Kobe, I. Barnaure, J. Galley, C. W. A. Pfirrmann, & F. Brunner, European Journal of Radiology (Vol. 129, p. 109044). Elsevier BV. https://doi.org/10.1016/j.ejrad.2020.109044
5) Alles, S. R. A., Cain, S. M., &Snutch, T. P. (2020). Pregabalin as a Pain Therapeutic: Beyond Calcium Channels [Review of Pregabalin as a Pain Therapeutic: Beyond Calcium Channels]. Frontiers in Cellular Neuroscience, 14. Frontiers Media. https://doi.org/10.3389/fncel.2020.00083
6) Amaya, S., Gónima-Valero, E., Mendoza, W. A. S., Sarmiento, D. A., & Andrade, M. J. (2022). Stellate ganglion block as an analgesic rescue alternative in phantom limb syndrome following an amputation:A case report. In S. Amaya, E. Gónima-Valero, W. A. S. Mendoza, D. A. Sarmiento, & M. J. Andrade, RevistaChilenade Anestesia (Vol. 51, Issue 5 Editorial Iku. https://doi.org/10.25237/revchilanestv5106071204
7) Aneja, R., Grover, R., Dhir, V., Shankar, S., & Kumar, A. (2006). Complex regional pain syndrome— management options. In R. Aneja, R. Grover, V. Dhir, S. Shankar, & A. Kumar, Indian Journal of Rheumatology (Vol. 1, Issue 3, p. 111). Medknow. https://doi.org/10.1016/s0973-3698(10)60203-8
8) Baidya, D. K., Kumar, A., Maitra, S., & Khanna, P. (2014). Clonidine for management of chronic pain: A brief review of the current evidences [Review of Clonidine for management of chronic pain: A brief review of the current evidences]. Saudi Journal of Anaesthesia, 8(1), 92. Medknow. https://doi.org/10.4103/1658-354x.125955
9) Baos, S., Rogers, C., Abbadi, R., Alzetani, A., Casali, G., Chauhan, N. R., Collett, L., Culliford, L., Jesus, S. E. de, Edwards, M., Goddard, N. J., Lamb, J., McKeon, H. E., Molyneux, M., Stokes, E. A., Wordsworth, S., Gibbison, B., &Pufulete, M. (2020). Effectiveness, cost-effectiveness and safety of gabapentin versus placebo as an adjunct to multimodal pain regimens in surgical patients: protocol of a placebo controlled randomised controlled trial with blinding (GAP study). In S. Baos, C. Rogers, R. Abbadi, A. Alzetani, G. Casali, N. R. Chauhan, L. Collett, L. Culliford, S. E. de Jesus, M. Edwards, N. J. Goddard, J. Lamb, H. E. McKeon, M. Molyneux, E. A. Stokes, S. Wordsworth, B.Gibbison, & M.Pufulete, BMJ Open (Vol. 10, Issue 11). BMJ. https://doi.org/10.1136/bmjopen-2020-041176
10) Bartl, R. (2022). Morbus Sudeck (CRPS). In R.

Bartl, Essentials. Springer Nature. https://doi.org/10.1007/978-3-662-66013-3

11) Bates, D., Schultheis, B. C., Hanes, M., Jolly,S., Chakravarthy, K., Deer, T. R., Levy, R. M., & Hunter, C. (2019). A Comprehensive Algorithm for Management of Neuropathic Pain [Review of A Comprehensive Algorithm for Management of Neuropathic Pain]. Pain Medicine, 20. Oxford University Press. https://doi.org/10.1093/pm/pnz075
12) Bell, R. F., &Kalso, E. A. (2018). Ketamine for pain management. https://journals.lww.com/01938936201810000-00007
13) Bhattacharya, A. K., Bhattacharyya, R., Ghosh, K. C., Das, S., Das, S., & Mondal, G. P. (2018). A Study of Post-stroke Pain in a Tertiary Care Hospital in Eastern India. In A. K. Bhattacharya, R. Bhattacharyya, K. C. Ghosh, S. Das, S. Das, & G. P. Mondal, Bengal Physician Journal (Vol. 5, Issue 3, p. 32). https://doi.org/10.5005/jp-journals-10070-5211
14) Birklein, F., & Dimova, V. (2017). Complex regional pain syndrome–up-to-date. In F. Birklein& V. Dimova, PAIN Reports (Vol. 2, Issue 6). Wolters Kluwer. ttps://doi.org/10.1097/pr9.0000000000000624
15) Bussa, M., Mascaro, A., Cuffaro, L., & Rinaldi, S. (2016). Adult Complex Regional Pain Syndrome Type I: A Narrative Review [Review of Adult Complex Regional Pain Syndrome Type I: A Narrative Review]. PM&R, 9(7), 707. Wiley. https://doi.org/10.1016/j.pmrj.2016.11.006
16) Campoy, L., Read, M., & Peralta, S. (2015). Canine and Feline Local Anesthetic and Analgesic Techniques (p. 827). https://doi.org/10.1002/9781119421375.ch45
17) Carr, E. S., De La Cerda, A., & Fiala, K. (2016). Complex Regional Pain Syndrome. https://www.tandfonline.com/doi/full/10.1080/08998280.2016.11929460
18) Castillo-Guzmán, S., Nava-Obregón, T. A., Palacios-Ríos, D., Estrada-Cortinas, J. Á., González-García, M., Mendez-Guerra, J. F., & González-Santiago, O. (2015). Complex regional pain syndrome (CRPS), a review [Review of Complex regional pain syndrome (CRPS), a review]. MedicinaUniversitaria, 17(67), 114. Elsevier BV. https://doi.org/10.1016/j.rmu.2015.03.003
19) Chapman, C. R., & Vierck, C. J. (2016). The Transition of Acute Postoperative Pain to Chronic Pain: An Integrative Overview of Research on Mechanisms [Review of The Transition of Acute Postoperative Pain to Chronic Pain: An Integrative Overview of Research on Mechanisms]. Journal of Pain, 18(4), 359. Elsevier BV. https://doi.org/10.1016/j.jpain.2016.11.004
20) Cheon, M., Kang, H. J., Hee, K., Yang, H. S., Han, E. J., & Yoo, J. (2021). Diagnostic Performance of Three-Phase Bone Scintigraphy and Digital Infrared Thermography Imaging for Chronic Post-Traumatic Complex Regional Pain Syndrome. In M. Cheon, H. J. Kang, K. Hee, H. S. Yang, E. J. Han, & J. Yoo, Diagnostics (Vol. 11, Issue 8, p. 1459). Multidisciplinary Digital Publishing Institute. https://doi.org/10.3390/diagnostics11081459
21) Chronic Pain Management. (2023). https://www.ncbi.nlm.nih.gov/books/NBK92054/
22) Cognitive Behavioral Therapy for Chronic Pain. (2023). https://www.va.gov/painmanagement/docs/cbtcp_therapist_manual.pdf
23) Complex Regional Pain Syndrome: An Update and Review of The Mechanisms. (2014). https://journals.lww.com/topicsinpainmanagement/citation/2014/10000/complex_regional_pain_syndrome__an_update_and.1.aspx
24) Costigan, M., Scholz, J., & Woolf, C. J. (2009). Neuropathic Pain: A Maladaptive Response of the Nervous System to Damage [Review of Neuropathic Pain: A Maladaptive Response of the Nervous System to Damage]. Annual Review of Neuroscience, 32(1), 1. Annual Reviews. https://doi.org/10.1146/annurev.neuro.051508.135531
25) DosSantos, M. F., Moura, B. de S., & DaSilva, A. F. (2017). Reward Circuitry Plasticity in

Pain Perception and Modulation [Review of Reward Circuitry Plasticity in Pain Perception and Modulation]. Frontiers in Pharmacology, 8. Frontiers Media. https://doi.org/10.3389/fphar.2017.00790

26) Drummond, P. D. (2010). Sensory Disturbances in Complex Regional Pain Syndrome: Clinical Observations, Autonomic Interactions, and Possible Mechanisms [Review of Sensory Disturbances in Complex Regional Pain Syndrome: Clinical Observations, Autonomic Interactions, and Possible Mechanisms]. Pain Medicine, 11(8), 1257. Oxford University Press. https://doi.org/10.1111/j.15264637.2010.00912.x

27) Eghbali, P., Becce, F., Goetti, P., Büchler, P., Pioletti, D. P., & Terrier, A. (2023). Glenohumeral Joint Force Prediction with Deep Learning. https://doi.org/10.2139/ssrn.4547277

28) FDA Approves Nasal Ketorolac for Short-Term Pain Management. (2010). In Topics in Pain Management (Vol. 26, Issue 2, p. 7). https://doi.org/10.1097/01.tpm.0000388148.18523.a9

29) Fuchs, P. N., Peng, Y. B., Boyette-Davis, J. A., &Uhelski, M. L. (2014). The anterior cingulate cortex and pain processing [Review of The anterior cingulate cortex and pain processing]. Frontiers in Integrative Neuroscience, 8. Frontiers Media. https://doi.org/10.3389/fnint.2014.00035

30) Galve Villa, M., Rittig-Rasmussen, B., Mikkelsen, L., Schear, M., Poulsen, A., &Groendahl. (2016). Complex regional

31) Gandhi, A. (2017). Stellate Ganglion Block: A Pediatric Case of Complex Regional Pain Syndrome. In A. Gandhi, Journal of Radiology Nursing (Vol. 36, Issue 2, p. 82). Elsevier BV. https://doi.org/10.1016/j.jradnu.2017.03.004

32) Ganu, S. S., & Pandya, Y. S. (2013). Complex Regional Pain Syndrome with Bilateral Lower Limb Involvement in an Adolescent. In S. S. Ganu & Y. S. Pandya, Journal of Musculoskeletal Pain (Vol. 21, Issue 1, p. 64). Informa. https://doi.org/10.3109/10582452.2012.762968

33) Greenwald, J. D., &Shafritz, K. M. (2018). An Integrative Neuroscience Framework for the Treatment of Chronic Pain: From Cellular Alterations to Behavior. In J. D. Greenwald & K. M. Shafritz, Frontiers in Integrative Neuroscience (Vol. 12). Frontiers Media. https://doi.org/10.3389/fnint.2018.00018

34) Harvey, M.-P., Maher-Bussières, S., Emery, E., Martel, M., Houde, F., Tousignant-Laflamme, Y., & Léonard, G. (2018). Evidence of motor system reorganization in complex regional pain syndrome type 1: A case report. In M.-P. Harvey, S. Maher-Bussières, E. Emery, M. Martel, F. Houde, Y. TousignantLaflamme, & G. Léonard, Canadian Journal of Pain (Vol. 2, Issue 1, p. 21). Taylor & Francis. https://doi.org/10.1080/24740527.2017.1422116

35) He, H., Hu, L., Tan, S., Tang, Y., Duan, M., Yao, D., Zhao, G., & Luo, C. (2023). Functional Changes of White Matter Are Related to Human Pain Sensitivity during Sustained Nociception. In H. He, L. Hu, S. Tan, Y. Tang, M. Duan, D. Yao, G. Zhao, & C. Luo, Bioengineering (Vol. 10, Issue 8, p. 988). Multidisciplinary Digital Publishing Institute. https://doi.org/10.3390/bioengineering10080988

36) Henry, D. E., Chiodo, A. E., & Yang, W.-B. (2011). Central Nervous System Reorganization in a Variety of Chronic Pain States: A Review [Review of Central Nervous System Reorganization in a Variety of Chronic Pain States: A Review]. PM&R, 3(12), 1116. https://doi.org/10.1016/j.pmrj.2011.05.018

37) Henry, J. L. (2014). Mechanisms of Peripheral Nerve Injury – What to Treat, When to Treat. In J. L. Henry, InTech eBooks. https://doi.org/10.5772/58421

38) Ishida, T., Dierks, T., Strik, W., & Morishima, Y. (2020). Converging Resting State Networks Unravels Potential Remote Effects of Transcranial Magnetic Stimulation for Major Depression. In T. Ishida, T. Dierks, W. Strik, & Y. Morishima, Frontiers in Psychiatry (Vol. 11). Frontiers

Pain Syndrome. https://www.ScienceWireLet.com/sci

Media. https://doi.org/10.3389/fpsyt.2020.00836

39) Islam, S., Nasim, M. A. A., Hossain, M. I., Ullah, Dr. M. A., Gupta, K. D., & Bhuiyan, M. M. H. (2023). Introduction of Medical Imaging Modalities (p. 1). https://doi.org/10.1007/978-3-031-47772-0_1

40) Jacob, A., Kaelin, D. L., Roach, A. R., Ziegler, C., & LaFaver, K. (2018). Motor Retraining (MoRe) for Functional Movement Disorders: Outcomes From a 1-Week Multidisciplinary Rehabilitation Program. In A. Jacob, D. L. Kaelin, A. R. Roach, C. Ziegler, & K. LaFaver, PM&R (Vol. 10, Issue 11, p. 1164). Wiley. https://doi.org/10.1016/j.pmrj.2018.05.011

41) Jang, S. H., & Seo, Y. S. (2020). Diagnosis of Complex Regional Pain Syndrome I Following Traumatic Axonal Injury of the Corticospinal Tract in a Patient with Mild Traumatic Brain Injury. In S. H. Jang & Y. S. Seo, Diagnostics (Vol. 10, Issue 2, p. 95). Multidisciplinary Digital Publishing Institute. https://doi.org/10.3390/diagnostics10020095

42) Jannic, A., Bernigaud, C., Brenaut, E., &Chosidow, O. (2018). Scabies Itch [Review of Scabies Itch]. Dermatologic Clinics, 36(3), 301. Elsevier BV. https://doi.org/10.1016/j.det.2018.02.009

43) Johanson, M. E. (2016). Rehabilitation After Surgical Reconstruction to Restore Function to the Upper Limb in Tetraplegia: A Changing Landscape. In M. E. Johanson, Archives of Physical Medicine and Rehabilitation (Vol. 97, Issue 6). Elsevier BV. https://doi.org/10.1016/j.apmr.2016.03.015

44) Kabani, R., & Brassard, A. (2014). Dermatological Findings in Early Detection of Complex Regional Pain Syndrome. In R. Kabani & A. Brassard, JAMA Dermatology (Vol. 150, Issue 6, p. 640). American Medical Association. https://doi.org/10.1001/jamadermatol.2013.7459

45) Kendroud, S., Fitzgerald, L. A., Murray, I., & Hanna, A. (2019). Physiology, Nociceptive Pathways. In S. Kendroud, L. A. Fitzgerald, I. Murray, & A. Hanna, StatPearls. https://europepmc.org/article/MED/29262164

46) Khan, S., Saud, S., Khan, I., Asif, M., Ismail, O., Salam, A., Yang, T., & Norville, K. J. (2018). Serotonin Syndrome Presenting with Concomitant Tramadol and Diphenhydramine Use: A Case Report of an Unlikely Side-Effect. In S. Khan, S. Saud, I. Khan, M. Asif, O. Ismail, A. Salam, T. Yang, & K. J. Norville, Cureus. Cureus, Inc. https://doi.org/10.7759/cureus.2421

47) Kortekaas, M. C., Niehof, S. P., Stolker, R. J., & Huygen, F. (2015). Pathophysiological Mechanisms Involved in Vasomotor Disturbances in Complex Regional Pain Syndrome and Implications for Therapy: A Review [Review of Pathophysiological Mechanisms Involved in Vasomotor Disturbances in Complex Regional Pain Syndrome and Implications for Therapy: A Review]. Pain Practice, 16(7), 905. Wiley. https://doi.org/10.1111/papr.12403

48) Krepska, A., Hastings, J., &Roodenburg, O. (2017). The metabolic and endocrine response to trauma. In A. Krepska, J. Hastings, & O. Roodenburg, Anaesthesia & intensive care medicine (Vol. 18, Issue 8, p. 414). Elsevier BV. https://doi.org/10.1016/j.mpaic.2017.05.007

49) Kutafina, E., Becker, S., & Namer, B. (2023). Measuring pain and nociception: Through the glasses of a computational scientist. Transdisciplinary overview of methods [Review of Measuring pain and nociception: Through the glasses of a computational scientist. Transdisciplinary overview of methods]. Frontiers in Network Physiology, 3. Frontiers Media. https://doi.org/10.3389/fnetp.2023.1099282

50) Kuttikat, A., Noreika, V., Chennu, S., Shenker, N., Bekinschtein, T. A., & Brown, C. (2017). Altered neurocognitive processing of tactile stimuli in patients with Complex Regional Pain Syndrome (CRPS). In A. Kuttikat, V. Noreika, S. Chennu, N. Shenker, T. A. Bekinschtein, & C. Brown, bioRxiv

(Cold Spring Harbor Laboratory). Cold Spring Harbor Laboratory. https://doi.org/10.1101/146613

51) Květňanský, R., Sabban, E. L., & Palkovits, M. (2009). Catecholaminergic Systems in Stress: Structural and Molecular Genetic Approaches [Review of Catecholaminergic Systems in Stress: Structural and Molecular Genetic Approaches]. Physiological Reviews, 89(2), 535. American Physiological Society. https://doi.org/10.1152/physrev.00042.2006

52) Lewiecki, E. M., Chastek, B., Sundquist, K., Williams, S. A., Weiss, R. J., Wang, Y., Fitzpatrick, L. A., & Curtis, J. R. (2020). Osteoporotic fracture trends in a population of US managed care enrollees from 2007 to 2017. In E. M. Lewiecki, B. Chastek, K. Sundquist, S. A. Williams, R. J. Weiss, Y. Wang, L. A. Fitzpatrick, & J. R. Curtis, Osteoporosis International (Vol. 31, Issue 7, p. 1299). Springer Science+Business Media. https://doi.org/10.1007/s00198-020-05334-y

53) Liang, D., Ren, Y., Huang, L., & Jin, S. (2022). A Study on THE Mechanism of Electroacupuncture to Alleviate Visceral Pain and NGF Expression. In D. Liang, Y. Ren, L. Huang, & S. Jin, Computational Intelligence and Neuroscience (Vol. 2022, p. 1). Hindawi Publishing Corporation. https://doi.org/10.1155/2022/3755439

54) Malkin, B., & Berridge, P. (2009). Guidance on maintaining personal hygiene in nail care. In B. Malkin & P. Berridge, Nursing Standard (Vol. 23, Issue 41, p. 35). Royal College of Nursing. https://doi.org/10.7748/ns2009.06.23.41.35.c7048

55) MariekeNiesters; Christian Martini; Albert Dahan. (2023). Ketamine for chronic pain: risks and benefits. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4014022/

56) Mathur, N., Jain, S. K., Kumar, N., Srivastava, A., Purohit, N., & Patni, A. (2014). Spinal Cord Injury: Scenario in an Indian State. In N. Mathur, S. K. Jain, N. Kumar, A. Srivastava, N. Purohit, & A. Patni, Spinal Cord (Vol. 53, Issue 5, p. 349). Springer Nature. https://doi.org/10.1038/sc.2014.153

57) McGonagle, D., Tan, A. L., & Benjamin, M. (2008). The Nail as a Musculoskeletal Appendage – Implications for an Improved Understanding of the Link between Psoriasis and Arthritis [Review of The Nail as a Musculoskeletal Appendage – Implications for an Improved Understanding of the Link between Psoriasis and Arthritis]. Dermatology, 218(2), 97. Karger Publishers. https://doi.org/10.1159/000182250

58) Mechanisms of Pain. (2023). https://www.ncbi.nlm.nih.gov/books/NBK32659/

59) Michael Costigan, Neural Plasticity Research Group, Harvard Medical School, 02129, Boston, Massachusetts, mcostigan@partners.org, Joachim Scholz, Neural Plasticity Research Group, Harvard Medical School, 02129, Boston, Massachusetts, scholz.joachim@mgh.harvard.edu, Clifford J Woolf, Neural Plasticity Research Group, Harvard Medical School, 02129, Boston, Massachusetts, cwoolf@partners.org. (2010). Neuropathic pain: a maladaptive response of the nervous system to damage. https://www.annualreviews.org/doi/10.1146/annurev.neuro.051508.135531

60) Musfiroh, I., Kartasasmita, R. E., Ibrahim, S., Muchtaridi, M., Hidayat, S., & Ikram, N. K. K. (2023). Stability Analysis of the Asiatic Acid-COX-2 Complex Using 100 ns Molecular Dynamic Simulations and Its Selectivity against COX-2 as a Potential Anti-Inflammatory Candidate. In I. Musfiroh, R. E.Kartasasmita, S. Ibrahim, M. Muchtaridi, S. Hidayat, & N. K. K. Ikram, Molecules (Vol. 28, Issue 9, p. 3762). Multidisciplinary Digital Publishing Institute. https://doi.org/10.3390/molecules28093762

61) NeuroLogic Examination Videos and Descriptions: Motor > Normal. (2020). https://neurologicexam.med.utah.edu/adult/html/motor_normal.html

62) Nuckolls, G. H., Kinnett, K., Dayanidhi, S., Domenighetti, A. A., Duong, T., Hathout, Y., Lawlor, M. W., Lee, S. S. M., Magnusson, S. P., McDonald, C. M., McNally, E. M., Miller, N., Olwin, B. B., Raghavan, P., Roberts, T. J., Rutkove, S. B., Sarwark, J. F., Senesac, C.,

Vogel, L. F., … Lieber, R. L. (2020). Conference report on contractures in musculoskeletal and neurological conditions [Review of Conference report on contractures in musculoskeletal and neurological conditions]. Muscle & Nerve, 61(6), 740. Wiley. https://doi.org/10.1002/mus.26845

63) Pagan-Rosado, R., Smith, B. J., Smither, F. C., Pingree, M. J., & D'Souza, R. S. (2023). Peripheral Nerve Stimulation for the Treatment of Phantom Limb Pain: A Case Series. In R. Pagan-Rosado, B. J. Smith, F. C. Smither, M. J. Pingree, & R. S. D'Souza, Case Reports in Anesthesiology (Vol. 2023, p. 1). Hindawi Publishing Corporation. https://doi.org/10.1155/2023/1558183

64) Pain Management. (2017). https://flylib.com/books/en/2.569.1.22/1/

65) Pain modulation in the spinal cord. (2023). https://www.ncbi.nlm.nih.gov/pmc/articles/PMC9513129/

66) Pappagallo, M., Breuer, B., Lin, H., Moberly, J. B., Tai, J., Noto, C., Sanchez, A., & Manfredi, P. L. (2013). A pilot trial of intravenous pamidronate for chronic low back pain. In M. Pappagallo, B. Breuer, H. Lin, J. B. Moberly, J. Tai, C. Noto, A. Sanchez, & P. L. Manfredi, Pain (Vol. 155, Issue 1, p. 108). Lippincott Williams & Wilkins. https://doi.org/10.1016/j.pain.2013.09.016

67) Park, B., Lee, J., Kim, H. J., Woo, C., & Park, H. (2020). A neuroimaging marker for predicting longitudinal changes in pain intensity of subacute back pain based on large-scale brain network interactions. In B. Park, J. Lee, H. J. Kim, C. Woo, & H. Park, Scientific Reports (Vol. 10, Issue 1). Nature Portfolio. https://doi.org/10.1038/s41598-020-74217-3

68) Pinho-Ribeiro, F. A., Verri, W. A., & Chiu, I. M. (2016). Nociceptor Sensory Neuron–Immune Interactions in Pain and Inflammation [Review of Nociceptor Sensory Neuron–Immune Interactions in Pain and Inflammation]. Trends in Immunology, 38(1), 5. Elsevier BV. https://doi.org/10.1016/j.it.2016.10.001

69) Rees, B., Borda, E., Nicolelis, M. A. L., & Yadav, A. P. (2022). Closed-loop spinal cord stimulation is superior in restoring locomotion in rodent models of Parkinson's Disease. In B. Rees, E. Borda, M. A. L. Nicolelis, & A. P.Yadav, bioRxiv (Cold Spring Harbor Laboratory). Cold Spring Harbor Laboratory. https://doi.org/10.1101/2022.09.19.508432

70) Resmini, G., Ratti, C., Canton, G., Murena, L., Moretti, A., & Iolascon, G. (2015). Treatment of complex regional pain syndrome [Review of Treatment of complex regional pain syndrome]. Clinical Cases in Mineral and Bone Metabolism. CIC Edizioni Internazionali. https://doi.org/10.11138/ccmbm/2015.12.3s.026

71) Reuben, D. B., Magasi, S., McCreath, H., Bohannon, R. W., Wang, Y., Bubela, D., Rymer, W. Z., Beaumont, J. L., Rine, R. M., Lai, J., & Gershon, R. (2013). Motor assessment using the NIH Toolbox. In D. B. Reuben, S. Magasi, H. McCreath, R. W. Bohannon, Y. Wang, D. Bubela, W. Z. Rymer, J. L. Beaumont, R. M. Rine, J. Lai, & R. Gershon, Neurology (Vol. 80, Issue 11). Lippincott Williams & Wilkins. https://doi.org/10.1212/wnl.0b013e3182872e01

72) Rockett, M. (2014). Diagnosis, mechanisms and treatment of complex regional pain syndrome [Review of Diagnosis, mechanisms and treatment of complex regional pain syndrome]. Current Opinion in Anaesthesiology, 27(5), 494. Lippincott Williams & Wilkins. https://doi.org/10.1097/aco.0000000000000114

73) Roghani, R. S., & Mansoor, S. (2012). Basics of Peripheral Nerve Injury Rehabilitation. In R. S.Roghani& S. Mansoor, InTech eBooks. https://doi.org/10.5772/39371

74) Rowbotham, M. C. (2006). Pharmacologic Management of Complex Regional Pain Syndrome [Review of Pharmacologic Management of Complex Regional Pain Syndrome]. Clinical Journal of Pain, 22(5), 425. Lippincott Williams & Wilkins. https://doi.org/10.1097/01.ajp.0000194281.74379.01

75) Saleem, M., & Naz, H. (2017). Analgesics: New Target and Sources. In M. Saleem & H.

Naz, InTech eBooks. https://doi.org/10.5772/67853

76) Saranita, J., Childs, D., & Saranita, A. D. (2008). Spinal Cord Stimulation in the Treatment of Complex Regional Pain Syndrome (CRPS) of the Lower Extremity: A Case Report. In J. Saranita, D. Childs, & A. D. Saranita, The Journal of Foot & Ankle Surgery (Vol. 48, Issue 1, p. 52). Elsevier BV. https://doi.org/10.1053/j.jfas.2008.10.003

77) Schilder, J. C. M., Niehof, S. P., Marinus, J., &Hilten, J. J. van. (2014). Diurnal and Nocturnal Skin Temperature Regulation in Chronic Complex Regional Pain Syndrome. In J. C. M. Schilder, S. P. Niehof, J. Marinus, & J. J. van Hilten, Journal of Pain (Vol. 16, Issue 3, p. 207). Elsevier BV. https://doi.org/10.1016/j.jpain.2014.11.012

78) Schneider, J., Patterson, M., & Jiménez, X. (2019). Beyond depression: Other uses for tricyclic antidepressants [Review of Beyond depression: Other uses for tricyclic antidepressants]. Cleveland Clinic Journal of Medicine, 86(12), 807. Cleveland Clinic. https://doi.org/10.3949/ccjm.86a.19005

79) Scholz, S., Lüthi, F., Wertli, M. M., & Brunner, F. (2019). Direct Health Care Cost and Work Incapacity Related to Complex Regional Pain Syndrome in Switzerland: A Retrospective Analysis from 2008 to 2015. In S. Scholz, F. Lüthi, M. M. Wertli, & F. Brunner, Pain Medicine (Vol. 20, Issue 8, p. 1559). Oxford University Press. https://doi.org/10.1093/pm/pnz030

80) Seidel, J., Bockhop, F., Mitkovski, M., Martin, S., Ronnenberg, A., Krueger-Burg, D., Schneider, K., Röhse, H., Wüstefeld, L., Cosi, F. G., Bröking, K., Schacht, A., & Ehrenreich, H. (2019). Vascular response to social cognitive performance measured by infrared thermography: A translational study from mouse to man. In J. Seidel, F. Bockhop, M. Mitkovski, S. Martin, A. Ronnenberg, D. Krueger-Burg, K. Schneider, H. Röhse, L. Wüstefeld, F. G. Cosi, K. Bröking, A. Schacht, & H. Ehrenreich, FASEB BioAdvances (Vol. 2, Issue 1, p. 18). Wiley. https://doi.org/10.1096/fba.2019-00085

81) Steinhoff, M., Schmelz, M., Szabó, I., &Oaklander, A. L. (2018). Clinical presentation, management, and pathophysiology of neuropathic itch [Review of Clinical presentation, management, and pathophysiology of neuropathic itch]. The Lancet Neurology, 17(8), 709. Elsevier BV. https://doi.org/10.1016/s1474-4422(18)30217-5

82) Subedi, M., Bajaj, S., Kumar, M. S., & YC, M. (2018). An overview of tramadol and its usage in pain management and future perspective [Review of An overview of tramadol and its usage in pain management and future perspective]. Biomedicine & Pharmacotherapy, 111, 443. Elsevier BV. https://doi.org/10.1016/j.biopha.2018.12.085

83) Tajerian, M., & Clark, J. D. (2015). New Concepts in Complex Regional Pain Syndrome [Review of New Concepts in Complex Regional Pain Syndrome]. Hand Clinics, 32(1), 41. Elsevier BV. https://doi.org/10.1016/j.hcl.2015.08.003

84) Tansey, E., Roe, S., & Johnson, C. (2014). The sympathetic release test: a test used to assess thermoregulation and autonomic control of blood flow. In E. Tansey, S. Roe, & C. Johnson, AJP Advances in Physiology Education (Vol. 38, Issue 1, p. 87). American Physical Society. https://doi.org/10.1152/advan.00095.2013

85) The Anatomy and Physiology of Pain. (2023). https://www.ncbi.nlm.nih.gov/books/NBK219252/ 86)The gate control theory of pain. (2023). https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1607474/

87) Topics in Pain Management CE Quiz. (2016). In Topics in Pain Management (Vol. 32, Issue 5, p. 11). https://doi.org/10.1097/01.tpm.0000510698.97902.fb

88) Tredici, K. D., & Braak, H. (2013). Idiopathic Parkinson's Disease: Staging an α-Synucleinopathy with a Predictable Pathoanatomy. https://www.ncbi.nlm.nih.gov/books/NBK6077/

89) Tsuda, M., Koga, K., Chen, T., & Zhuo, M. (2017). Neuronal and microglial mechanisms for neuropathic pain in the

spinal dorsal horn and anterior cingulate cortex [Review of Neuronal and microglial mechanisms for neuropathic pain in the spinal dorsal horn and anterior cingulate cortex]. Journal of Neurochemistry, 141(4), 486. Wiley. https://doi.org/10.1111/jnc.14001

90) Vahedifar, P., &Kamrava, E. (2015). Complex Regional Pain Syndrome Types I and II. In P. Vahedifar& E. Kamrava, Springer eBooks (p. 17). Springer Nature. https://doi.org/10.1007/978-3-319-21587-7_3

91) Volcheck, M. M., Graham, S. M., Fleming, K. C., Mohabbat, A. B., & Luedtke, C. A. (2023). Central sensitization, chronic pain, and other symptoms: Better understanding, better management [Review of Central sensitization, chronic pain, and other symptoms: Better understanding, better management]. Cleveland Clinic Journal of Medicine, 90(4), 245. Cleveland Clinic. https://doi.org/10.3949/ccjm.90a.22019

92) Yang, S., & Chang, M. C. (2019). Chronic Pain: Structural and Functional Changes in Brain Structures and Associated Negative Affective States [Review of Chronic Pain: Structural and Functional Changes in Brain Structures and Associated Negative Affective States]. International Journal of Molecular Sciences, 20(13), 3130. Multidisciplinary Digital Publishing Institute. https://doi.org/10.3390/ijms20133130

93) Yousefshahi, F., Predescu, O., & Asenjo, J. F. (2017). The Efficacy of Systemic Lidocaine in the Management of Chronic Pain: A Literature Review [Review of The Efficacy of Systemic Lidocaine in the Management of Chronic Pain: A Literature Review]. Anesthesiology and Pain Medicine, 7(3). Kowsar Medical (Netherlands). https://doi.org/10.5812/aapm.44732